Journal of Ayurveda and Integrative Medicine 9 (2018) 126-130

Contents lists available at ScienceDirect

Journal of Ayurveda and Integrative Medicine

journal homepage: http://elsevier.com/locate/jaim



AYURVEDA

TRANSDISCIPLINARY

An exploratory clinical study to determine the utility of heart rate variability analysis in the assessment of *dosha* imbalance



J-AIN



P. Ram Manohar ^{a, *}, Oleg Sorokin ^{b, c}, James Chacko ^d, Vasudevan Nampoothiri ^d

^a Amrita Centre for Advanced Research in Ayurveda, Amrita University, Clappana P.O., Vallikkavu, Kollam, Kerala 690525, India

^b National Ayurvedic Medical Association, Russian Federation

^c Biokvant LLC, 1st Parkhomenko Alley, d. 14, Office 7, Novosibirsk 630108, Russian Federation

^d Amrita School of Ayurveda, Amrita University, Clappana P.O., Vallikkavu, Kollam, Kerala 690525, India

ARTICLE INFO

Article history: Received 3 March 2017 Received in revised form 31 May 2017 Accepted 16 June 2017 Available online 19 December 2017

Keywords: Heart rate variability Ayurveda Spectral analysis

ABSTRACT

The present study is a comparison of the data of spectral analysis of heart rate variability with clinical evaluation of pathological state of *doshas*. The calculated cardiointervalography values are combined into three integral indexes, which according to the authors' opinion reflect the influence on heart rhythm of *vata, pitta* and *kapha*, the regulation systems of the body known as *doshas* in Ayurveda. Seven gross *dosha* imbalances were assessed to test the agreement between the two methods in this study. Heart Rate Variability (HRV) spectral data was collected from 42 participants to make the comparison with the clinical assessment of *dosha* imbalance. Clinical method of *dosha* assessment and method of calculating integral indexes by cardiointervalography data showed substantial agreement by Kappa coefficient statistic (k = 0.78) in assessment of gross *dosha* imbalance. The results of the data generated from this pilot study warrant further studies to rigorously validate the algorithms of HRV analysis in understanding *dosha* imbalance in Ayurvedic clinical practice and research settings.

© 2017 Transdisciplinary University, Bangalore and World Ayurveda Foundation. Publishing Services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

1. Introduction

Ayurveda explains homeodynamics and health in terms of balance of the three *doshas* (regulatory systems), whereas disease is the outcome of the failure to maintain homeodynamics and is understood in terms of the imbalance of *doshas*. In order to formulate a specific treatment protocol, an Ayurvedic physician has to determine the exact nature of imbalance of the *doshas* and the specific substratum in which the imbalance manifests. To a great extent, imbalance of the *doshas* can be understood by careful analysis of clinical symptoms and signs. However, in many instances, it can be challenging to compute the imbalance in terms of the dominance and relative increase or decrease of the *doshas*. In ancient times, Ayurvedic physicians developed protocols for detailed examination of tongue, eyes, urine, faeces, skin and so on to measure imbalance of *doshas*. Subtle techniques like pulse

* Corresponding author.

E-mail: dr.rammanohar@ayurveda.amrita.edu

Peer review under responsibility of Transdisciplinary University, Bangalore.

diagnosis were introduced later on and eventually became the gold standard for diagnosing *dosha* imbalance.

1.1. HRV and imbalance of the doshas?

HRV is the result of the impact of the autonomous regulation on the heartbeat. Doshas are also autonomous regulators of body physiology. It is well known that 90% of our responses to the challenges of the external environment are determined by the autonomous nervous system. Travis and Wallace have discussed about the possible correlations between autonomous nervous system and the doshas [1]. Tyagi and Cohen have reviewed evidence indicating that physiological and psychological stress disrupts autonomic balance, which has long-term implications in a wide range of mental and physical illnesses [2]. Several studies have demonstrated that HRV is a useful tool in assessing pathology as well as treatment outcomes in many diseases. Masel et al. showed that HRV could be a surrogate marker for alleviation of cancer through pain and could also detect pain without active participation of the patients [3]. Kim et al. have demonstrated that perioperative HRV correlated with pre-operative depressed mood

http://dx.doi.org/10.1016/j.jaim.2017.06.008

^{0975-9476/© 2017} Transdisciplinary University, Bangalore and World Ayurveda Foundation. Publishing Services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

in patients with hepatic cancer [4]. Koszewicz et al. profiled autonomic dysfunctions in patients with primary brain tumor revealing sympathetic hyperactivity through HRV analysis [5]. Zhou et al. have reviewed the role of HRV in the prediction of survival in patients with cancer [6]. Taffe et al. have assessed post prandial HRV spectral analysis to differentiate overweight from normal weight adults [7]. Gupta et al. concluded that a significantly raised central vagal outflow and a concomitant significantly low central sympathetic efferent could be appreciated in asymptomatic asthmatic patients as compared to that in the control group by HRV spectral data analysis [8]. These studies indicate that HRV can vary in different diseases and also in specific stages of different diseases, which is also the case with the Tridoshas. This study aims to generate pilot data that will help in understanding the correlation between dosha imbalance and HRV patterns in patients diagnosed with different clinical conditions.

1.2. Previous studies exploring the correlation between HRV and doshas

Recently, Harupjit Singh from the Electrical and Instrumentation Engineering Department Thapar University, Patiala explored the relationship between three Ayurvedic *doshas* and HRV frequency bands by conducting a pilot study for his masters thesis [11]. His study involved only twenty-five patients and it achieved classification accuracy of 53% in case of the *vata dosha* and 70% in case of the *pitta dosha*.

2. Materials and methods

2.1. Study description

For the purpose of this study, forty-two patients with different Ayurvedic constitutions and pathology were randomly selected. All patients were attending the Amrita School of Ayurveda Hospital and signed a voluntary informed agreement to participate in the study.

Criteria for exclusion from the study included: medications for cardiac ailments influencing chronotropy of sinus node; multiple heart rhythm failures; divergence in data in verification of pathological *dosha* dominance between checklist of *dosha* dominance symptoms and expert assessment; HRV recording with many artefacts.

A checklist of *dosha* dominance symptoms compiled from the classical texts was used to make an exhaustive listing of the clinical symptomatology of the patient. The patients were then seen by a team of clinical experts who assessed the *dosha* dominance by performing an independent clinical assessment of the patient. Eleven participants with discrepancy in *dosha* dominance based on checklist and clinical assessment were excluded from the study and forty-two participants with congruence in *dosha* dominance based on checklist and clinical assessment were included for HRV analysis.

Researchers who filled the *dosha* dominance checklist, the clinical experts and the experts who recorded HRV spectral data were mutually blinded from the results of the independent assessments.

A VedaPulse hardware and software kit was used (manufactured by Biokvant LLC, Russia) for objective quantitative assessment of pathological *dosha* dominance using algorithms of HRV analysis. These algorithms were developed with inputs from Ayurvedic professionals with expertise in pulse diagnosis as well as pilot studies in a number of people.

Analysis of pathological *dosha* dominance included sequential actions, including: registration of biopotentials of the work of the heart for 5 min by placing electrodes on the wrists, software filtration of the signal with further receiving of HRV periodogram and analysis of cardiointervalogram using methods of spectral data analysis [9].

Frequency analysis of cardiointervalogram was done using discrete Fourier transform with overlapping. Periodogram was divided into three equal parts and for each of them a Fourier spectrum was calculated, after which these spectrums were averaged (see Fig. 2). Using the Fourier analysis, we estimate three spectral intervals: VLF — interval of very low frequencies (blue color); LF — interval of low frequencies (red color); HF — interval of high frequencies (green color). Horizontal axis shows frequencies in Hz, vertical axis shows spectrum power in ms2.

Practically the quantitative calculation of the *dosha* balance was done the following way: we calculated absolute values of the spectrum and rated them using 10-grade system of measurement, in which case the values of 2–4 points were considered a relatively standard interval which roughly corresponds with the average population value of the norm by absolute values of spectrum (see Fig. 3). If the normalized values of *doshas* in points were higher than the indicated interval and spectrums in VLF and LF intervals were dominating, it corresponded with–*vata-pitta*. If



Fig. 1. Cardiac cycle duration variations.



Fig. 2. VLF, LF and HF spectrums.

VLF and HF spectrums dominated, it corresponded with *vata-kapha*. If LF and HF spectrums dominated, it corresponded with *pitta-kapha*. If all three spectrums had a rising tendency, it corresponded with VPK. In Fig. 3, Yellow bar (Factor "V") – total value of *vata dosha*; red bar (Factor P) – total value of *pitta dosha*; blue bar (Factor K) – total value of *kapha*. Corridor of norm 2–4 is highlighted in green.

To detect frequency intervals, we use the following frames:

For VLF – (Time = 25-300 s, frequency = 0.04-0.0033 Hz) For LF – (Time = 6.6-25 s, frequency = 0.15-0.04 Hz) For HF – (Time = 2.5-6.6 s, frequency = 0.4-0.15 Hz)

A group of scientists under the leadership of Dr. Sorokin began to do comparative analysis of the HRV data with the clinical analysis of the *dosha* balance starting from 2006, the results of which were published. Later, similar algorithms for calculating *doshas* by HRV were offered in the works of Kelkar et al. [10] and Harupjit Singh [11], where they showed the connection of these intervals with *doshas* on the sample of several hundred participants. The difference of our work is that the power of spectrum in intervals was rated using a 10-grade system, which simplified the comparison with the scoring system of *dosha* calculation by clinical method. Normalizing of the absolute spectral power to a 10-grade scale is necessary to create relative values of the secondary indices – factor V, factor P, factor K, which reflect the *dosha* balance. The use of such normalizing eases the perception of the balance of these factors between them, which is hard when comparing the absolute values of the specter. This normalizing does not change the mathematic approach to the calculation, it only facilitates better visualization of the data.

2.2. Scales to assess dosha dominance

There are simple to complex scales to assess *dosha* dominance and simplest scale categorizing *dosha* dominance in seven categories was chosen for comparing diagnostic agreement in this study. The most complex scale of *dosha* dominance has sixty-three categories, which take into account the permutations and combinations of the relative increase and decrease of the *doshas*.

These seven categories cover the single, dual (samsarga) and triple (sannipāta) combinations of *doshas* that are routinely assessed in clinical practice. At the next level of complexity, distinction is made between *vata-pitta* and *pitta-vata*, as well as between *vata-kapha* and *kapha-vata* and between *kapha-pitta* and *pitta-kapha*. At even higher levels of complexity, the relative increase and decrease of the *doshas* are estimated. For example, *vata* increased, *pitta* decreased, *kapha* normal and so on and so forth.

Dosha assessment was made by clinical examination of study participants by experts, which was then compared with HRV spectral data. Algorithms based on HRV data were used to assess



Fig. 3. Balance of the doshas.

dosha dominance from ECG signals. By a mathematical analysis of high frequency, low frequency and very low frequency signals from the ECG readings, the dominant influence of the *doshas* on the regulatory systems of the body were calculated. Figs. 1–3 illustrate the capture of ECG signals and interpretations in terms of the *doshic* dominance by the HRV analysis. In Fig. 1, the upper graph shows the calculation of cardiac cycle duration in milliseconds via measuring the RR intervals of ECG. The lower graph shows the dependence of cardiac cycle duration (vertical axe, in ms) from the numerical order of cardiointerval (horizontal axe).

3. Results

Table 1 gives the genderwise distribution of the study participants and it can be seen that nearly two-third were female. Table 1 gives the age group wise distribution of the study participants revealing that two-third were above the age of 50 and the maximum number of participants were in the age group of 61–70. The numbers in the tables indicate the frequency of participants in a specific category.

Interestingly, both HRV analysis and clinical assessment did not identify extreme *pitta* or *kapha* imbalances. There were very few participants for the extreme *vata*, *kapha-pitta* and *Tridosha* imbalances. Majority of the participants were identified as presenting either *vata-pitta* or *vata-kapha* imbalances (see Table 4).

Table 3 lists the clinical diagnosis of the study participants who participated in the study.

Out of the forty-two participants who completed the study, twenty-seven (64.28%) showed complete agreement and ten (23.80%) showed partial agreement in assessment made by HRV analysis and clinical assessment. There was complete disagreement in five participants (11.90%). In the case of partial agreement, there was a mismatch in the assessment of the second *dosha* (see Table 4).

In order to assess the diagnostic agreement between the HRV analysis and the clinical assessment, the Kappa co-efficient was calculated using seven categories. The results are tabulated in Table 2. The values in the diagonal of the table indicate the number of participants classified by HRV analysis and clinical assessment into the same category. Values in other cells indicate number of participants in which disagreement was observed.

Kappa co-efficient is a statistical measure to assess inter-rater agreement more accurately than a simple percentage analysis. It is helpful to assess diagnostic tests that involve some degree of subjective interpretation by the observers. Its reliability lies in its ability to take into account the agreement occurring by chance. A Kappa score of 1 indicates perfect agreement and a score of 0 indicates complete disagreement. Even a score of 0.41–0.60 indicates substantial agreement, while a score of 0.61–0.80 indicates substantial agreement. A score of 0.81–0.99 indicates perfect agreement [12].

Analysis of the data gave Kappa statistic score of 0.78, which means substantial agreement between the HRV analysis and the clinical method of assessing *doshic* imbalances in pathological conditions.

ladie I	
Age and Gender wise	distribution of patients.

Age group	21-30	31-40	41-50	51-60	61-70	71-80	81-90	Total
Male	1	2	2	3	6	1	1	16
Female	2	4	3	5	9	3	0	26
Total	3	6	5	8	15	4	1	42

Table 2

Dosha vikriti categories of study participants.

	HRV analysis	Clinical assessment		
V	5	2		
Р	0	0		
K	0	0		
VP	15	17		
VK	17	21		
KP	3	1		
VPK	2	1		

Table 3

The clinical diagnosis of the study participants who participated in the study.

Allopathic diagnosis (number of participants)	Ayurvedic diagnosis
Brachial neuralgia (1)	Apabahukam
Cervical spondylosis (1)	Manyasthambham
Diabetes melittus (1)	Prameha
Gouty arthritis (2)	Vataraktam
Ligament tear (1)	Marmabhighatam
Low back ache (2)	Katigraha
Lumbar spondylosis (3)	Katigraha
Obesity (1)	Sthoulya
Osteoarthritis (11)	Sandhigata vata
Partial paralysis (1)	Vatavyadhi
Peripheral neuropathy (1)	Padasupthi
Psoriasis (1)	Sidhma kushtam
Rheumatoid arthritis (4)	Amavata
Sciatica (11)	Gridhrasi
Sinusitis (1)	Peenasa

Table 4

Diagnostic agreement between HRV analysis and clinical assessment of dosha.

Clinical method	HRV analysis						
	v	Р	К	VP	VK	KP	VPK
V	2	0	0	0	0	0	0
Р	0	0	0	0	0	0	0
K	0	0	0	0	0	0	0
VP	1	0	0	11	2	2	1
VK	2	0	0	4	14	1	1
KP	0	0	0	0	1	0	0
VPK	0	0	0	0	0	0	0

4. Discussion

In the context of Ayurveda, *doshas* are the functional parameters which auto-regulate the physiology and therefore can have an impact on the heart rhythm.

Therefore, if there is a continuity between the formation of the pulse wave and the work of cardiac pacemakers, then instrumental analysis of HRV brings an important input to the understanding of physiological regulation processes, which connects disease development with pathological *dosha* dominance.

Modern technologies of biosignal registration and mathematical analysis refracted through the prism of empirical knowledge of Ayurvedic medicine allowed to develop an algorithm for assessing psychophysiological constitution and pathological *dosha* dominance. Analyzing spectral and temporal characteristics of HRV, a quantitative assessment system was developed to assess pathological *dosha* dominance, which was manifested in the VedaPulse hardware and software kit. This allowed building a connection between capabilities of modern electrophysiology and Ayurvedic medicine. The spectral analysis was carried out on the basis of the assumption that VLF, LF and HF grossly reflect *vata*, *pitta* and *kapha* respectively. Assessment of the *dosha* imbalance is of cardinal importance in Ayurvedic diagnosis and treatment of diseases. However, the assessment is highly subjective and there is no validated instrument available for this purpose. There is a high inter-rater variability amongst experts in the field of Ayurveda. For this reason, only the gross *dosha* imbalances were selected for comparison in this study.

Examination of pulse is expected to give deeper insights into the state of *doshic* imbalance than mere clinical assessment. Therefore, clinical assessment is not the preferred standard to test the accuracy of HRV based algorithms in assessment of *dosha* imbalance. The advantage is that the clinical method of *dosha* assessment based on symptom analysis is more objective than pulse analysis by experts. Series of studies comparing HRV spectral analysis with clinical assessment of *dosha* dynamics could potentially help in identifying and characterizing HRV signatures that can predict imbalance of the *dosha*.

The present study has several limitations. HRV based algorithms used in the study to assess *dosha* imbalance are still in the early stages of testing and validation. Clinical assessment of *doshas* is not considered to be as accurate as assessment by pulse examination. An objective questionnaire for clinical assessment of *dosha* imbalance is not yet developed and validated. There is inter-rater variability amongst experts in clinical assessment of *doshas*. Matched control and adoption of non-linear mathematics and time domain analysis will be undertaken in future studies. Taking these limitations into consideration, only gross imbalance of *doshas* with consensus of experts were included for comparison with HRV based algorithms.

A larger multicentric study extending over a longer period that accounts for diagnostic error based on treatment outcomes will be able to make more accurate assessment of the scope of HRV analysis to determine imbalance of *doshas*.

5. Conclusions

Preliminary study to assess the diagnostic agreement in assessment of gross *dosha* imbalance by clinical method and HRV analysis showed substantial agreement as revealed by the Kappa score of 0.78.

The present study demonstrates the scope and relevance of further studies to validate the utility of the HRV in assessment of complex *dosha* imbalances and other parameters like *dhatu* imbalance.

Use of validated questionnaires for *dosha* analysis is recommended for future studies to assess diagnostic accuracy of HRV analysis. HRV analysis is not expected to replace the assessment of the physician by palpation and clinical examination. On the other hand, it could serve as a useful tool for research purposes and as an aid in the clinical practice of Ayurveda.

Sources of funding

The authors acknowledge the financial support received from Biokvant LLC, Russia for the conduct of the study.

Conflict of interest

The second author of the paper, Dr Oleg Sorokin is also sponsor of the study and the developer of the HRV analysis hardware device and the software tool used in the study (VedaPulse).

Acknowledgments

The authors are thankful to the Research Dean of Amrita Vishwa Vidyapeetham University and the Medical Director of Amrita School of Ayurveda for providing facilities to conduct the study. We also acknowledge the contributions of Dr. Rajalakshmi, Research Staff, Dr. Gopikrishna and Dr. Krishnaprabha, PG Scholars, Athira CP and Aparna S, Internees and Dr. Devipriya, Faculty of Amrita School of Ayurveda for providing assistance in the conduct of the study.

References

- Travis FT, Wallace RK. Doşa brain-types: a neural model of individual differences. J Ayurveda Integr Med 2015;6(4):280–5.
- [2] Tyagi A, Cohen M. Yoga and heart rate variability: a comprehensive review of the literature. Int J Yoga 2016;9(2):97–113.
- [3] Masel E, Huber P, Engler T, Herbert Watzke HH. Heart rate variability during treatment of breakthrough pain in patients with advanced cancer: a pilot study. J Pain Res 2016 Dec;9:1215–20.
- [4] Kim E-H, Park J-H, Lee SM, Gwak M-S, Kim G-S, Kim M-H. Preoperative depressed mood and perioperative heart rate variability in patients with hepatic cancer. J Clin Anesth 2016 Dec;35:332–8.
- [5] Koszewicz M, Michalak S, Bilinska M, Budrewicz S, Zaborowski M, Slotwinski K, et al. Profile of autonomic dysfunctions in patients with primary brain tumor and possible autoimmunity. Clin Neurol Neurosurg 2016 Dec;151:51–4.
- [6] Zhou X, Ma Z, Zhang L, Zhou S, Wang J, Wang B, et al. Heart rate variability in the prediction of survival in patients with cancer: a systematic review and meta-analysis. J Psychosom Res 2016 Oct;89:20–5.
- [7] Taffe L, Stancil K, Bond V, Pemminati S, Gorantla VR, Kadur K, et al. Differentiation of overweight from normal weight young adults by postprandial heart rate variability and systolic blood pressure. J Clin Diagnostic Res 2016 Aug;10(8):CC01–6.
- [8] Gupta J, Dube A, Singh V, Gupta RC. Spectral analysis of heart rate variability in bronchial asthma patients. Indian J Physiol Pharmacol 2012;56(4):330–6.
- [9] Sorokin OV, Subotyalov MA. Comparative medical research of VedaPulse hardware and software kit // Health maintaining activities in educational system: theory and practice. In: Material of the all-Russian scientific and practical conference with international presence, October 26–27, 2011 Kemerovo, Russia; 2011. p. 225–30.
- [10] Kelkar P, Karamchandani S, Jinda S. Identifying tridosha for disease characterisation in morphology of an IPG pulse waveform. In: Proceedings of conference on advance applications in physiological variability, BARC; 2010.
- [11] Singh Harupjit. Relationship between three ayurvedic doshas and heart rate variability frequency bands: a pilot study. Patiala: Thapar University; 2016.
- [12] Anthony JV, Joanne MG. Understanding interobserver agreement: the kappa statistic. Fam Med 2005 May;37(5):360–3.