

RESEARCH

Open Access



Regional trade of medicinal plants has facilitated the retention of traditional knowledge: case study in Gilgit-Baltistan Pakistan

Muhammad Asad Salim^{1,2}, Sailesh Ranjitkar^{1,3}, Robbie Hart⁴, Tika Khan⁵, Sajid Ali⁵, Chandni Kiran⁵, Asma Parveen⁵, Zahra Batool⁵, Shanila Bano⁵ and Jianchu Xu^{1,3*} 

Abstract

Background: The ethnic groups in Gilgit-Baltistan have been utilizing local resources in their centuries-old traditional healing system. Most tribes within these ethnic groups still rely on traditional healing systems. We aim to understand the current status, uses, and abundance of medicinal plants, associated traditional knowledge, and trade.

Materials and methods: The study incorporated over 300 local community members (70% men and 30% women) in focused group discussions, semi-structured interviews, and homework assignments for 8th to 12th grade students to document traditional knowledge (TK) in six districts in Northeast Pakistan. We calculated various indices such as informant consensus factor, use value, relative frequency of citation, and CoKriging. These indices, along with repetitively used medicinal plants, were used to analyze differences in studied locations.

Results: Most of the community members still rely on traditional medication in the study areas. However, we found the highest number of medicinal plants used in Skardu and Gilgit compared to other districts and these two districts also represent trade centers and a highly populated area regarding medicinal plants. Results indicate connection amongst the surveyed villages signifying mixing of knowledge from different sources, with certain areas more influenced by traditional Chinese medicine and others more by Ayurveda and Unani.

Conclusion: TK is mostly retained with elder community members; however, those directly linked with market value chain retain rich knowledge on traditional use of the medicinal plants from the region. Major trade centers in the region also coincide with a high density of medicinal plant occurrence, knowledge, and higher utilization. Therefore, with the increasing trade in medicinal plant in the region, there is potential for rejuvenation of this knowledge and of plant use in the region.

Keywords: Ethnomedicine, Ethnoecology, Medicinal plants, Traditional knowledge, Trade in medicinal plants, Gilgit-Baltistan

* Correspondence: jxu@mail.kib.ac.cn

¹Key Laboratory for Plant Diversity and Biogeography of East Asia, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, Yunnan, China

³World Agroforestry Centre (ICRAF), East and Central Asia Office, Kunming 650201, Yunnan, China

Full list of author information is available at the end of the article



© The Author(s). 2019 **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

Introduction

Gilgit-Baltistan, the northeast mountainous region of Pakistan, is outstanding for its rich ethnic diversity [1–3]. This region is situated in a strategic geographical location that is important as a part of the ancient silk route and its position along the China-Pakistan Economic Corridor [3–5]. For centuries, knowledge exchange has occurred between indigenous dwellers and migrants and peddlers from the southern, northern, and western parts of Asia. Besides, the region is one of the important plant areas in the Himalaya, Karakoram, and Hindukush (KKH) landscape [6–12]. In the epoch of Anthropocene, as for other parts of the world, this region also experienced problems due to human population growth and associated land use transformation that severely affected both important species and the wider ecosystem [13–15]. Still natural resources, especially plant diversity, are very significant for ethnic communities in this mountainous landscape [10, 14, 16, 17]. Plants are the source of energy and food, a building material for houses, and a main component of the health care system as folklore medicine [18–27].

There are about 600 species of flowering plants in Pakistan that are utilized as medicinal plants and around 500 of these have global significance and studies available [9, 16, 28]. Around 50,000 traditional healers and informal Pansaris (retailers) are registered in Pakistan who frequently utilize and sell 400–600 plants species for their medicinal, cultural, traditional, and spiritual benefits [29–31]. Gilgit-Baltistan, with above 300 reported species of MAPs, is a hotspot for medicinal plants and their utilization in Pakistan [9, 12–14, 16, 27, 30, 32–41]. With seven districts and an ideal mountainous landscape, this region is naturally suitable for high-value medicinal plants. The local communities have been utilizing this resource for many generations [9, 12, 15, 33, 36, 42–45]. Several researchers have previously documented traditionally used medicinal plant from different districts in this region. Shadayi and Gulshan (2012) and Shadayi et al. (2014) in Ghizer district; Khan et al. (2013) and Akhtar et al. (2016) in Hunza; Bano et al. (2014) in Skardu; Khan and Khatoon (2007) and Fahad and Bano (2012) in Gilgit; Abbas et al. (2016) in Baltistan and the contributions of Hussain et al. (2011), Khan et al. (2011), Qureshi et al. (2006), and Ali et al. (2017) are important and noteworthy for the documentation of plant species used locally for medicinal purposes besides reporting on the modes of their uses and diseases targeted/cured through traditional herbal practitioners.

In addition to the utilization of medicinal plants in the traditional healing system, trade in herbal raw material and product is not new in the area. As this region serves as an ancient trade route that connects south Asia with China, Central Asia, and West Asia; trade in medicinal plants and

exchange of traditional healing knowledge is very likely. For instance, archeological studies reveal the influence of cultural incursions from the Indian subcontinent, China, Scythia (Eurasia), Transoxiana (Uzbekistan, Tajikistan, southern Kyrgyzstan, and southwest Kazakhstan), and Ancient Greece, amongst others on traditional medicinal system [3, 46]. Before the introduction of Islam, the communities in Gilgit-Baltistan were predominantly practicing Buddhism [3, 46–50]. The region is recognized for its contributions towards survival and expansion of Ayurveda during the British regime [51]. Hakim Ajmal Khan is a famous Indian physician who worked for the revival of Ayurveda and Unani systems during the British era by establishing an Ayurveda and Unani medical college and a pharmaceutical company besides continuing with his own clinical practice of the systems [52]. Before the region completely came under Dogra Raj of the Kashmir State, Hunza, Gilgit, Nagar, and Ghizer mainly remained under Chinese influence, whereas Skardu, Astore, Ghanche, and Diamer remained under Tibetan influence [3, 46]. Recently, with development of the Karakoram Highway (KKH) and China-Pakistan Economic Corridor (CPEC) that follow the ancient Silk Routes [3, 50] in the region, this area has become important junction of trade.

Our research provides the first comparative study for six districts of Gilgit-Baltistan where the influence of Chinese, Ayurveda, Unani, and Tibetan healing systems on folk knowledge has been observed. Traditional knowledge of medicinal plants is often socially integrated through communal learning and intercultural exchange. Medicinal plants in mountainous terrain are known for their distribution in elevation corridors or endemism to a particular locality. We therefore aim to explore the richness of traditional knowledge of medicinal plants, their uses, distribution, and trade in the mountainous region of Gilgit-Baltistan, learn about the knowledge exchange between the old and young generation as well as amongst different communities and localities of the region, and how medicinal systems like Chinese, Ayurveda, Tibetan, and Unani influenced the use of traditional medicine system in the region. We also explore the possible factors behind the general decline in knowledge about medicinal plants, yet the continued use of traditional medicine for treatment of different diseases and how the current markets and market players supplement this phenomenon. We take an inventory of these plants, their use value, and local importance at a regional level and current markets. We also look at how these factors and the influence of different medicinal systems in the region compliment the transfer of knowledge to younger generations and across different ethnicities and locations in the region.

Materials and methods

Study area

Gilgit-Baltistan is located in the far north of Pakistan, with Afghanistan to the north and west, China to the north and east, and India to the south [8]. The seven districts are spread over an area of 72,496 km². The region is rugged and mountainous, located amongst three of the highest mountain ranges—the Himalayas, the Karakoram, and the Hindukush (HKH) and home to the largest number of glaciers outside the polar region [36, 53, 54].


The study was focused on Gilgit, Hunza, Nagar, Ghizer, Skardu, and Astore Districts of Gilgit-Baltistan. The residents are divided into sub-groups based on their origin as well as their ethno-linguistic clustering (Fig. 1). Yashkun, Sheen/Shinaki, Wakhi (those who migrated from Wakhan), Burushos, Dom, Brokpa, and Balti are the main tribes of the area; some Kashmiris, Kohistani, Mongols, Mughals, Rajas, Pathans, Gujar, Soniwal, Mon, Hor, and Kashgari also reside here [35, 55–60]. The majority of Gilgit-Baltistan is sparsely populated with these tribes, but Ghizer is dominated by Burushos; Gilgit, Hunza, and Nagar have Burushos, Sheen,

and Yashkun; Diamer and Astore are majorly populated by Sheen, Yashkun, and Kohistani communities; and Skardu and Ghanche are predominantly Mon, Hor, and Mongols [60, 61]. The languages spoken by Burushos, Sheen, and Yashkun are Shina, Burushaski, Wakhi, and Khowar (only Ghizer and parts of Hunza) while the Mongols, Mon, and Hor speak Balti [9, 38, 60, 62].

Communities in Gilgit-Baltistan are dependent on agricultural resources and live close to the forest area (Fig. 1). Land cover changes, lack of resource management and sustainable harvesting policies, and political interests at massive scale in the HKH and Pamir mountain ranges have severe and long-lasting impacts on the region [63–66]. Medicinal plants and traditional medication have been used for generations for curing different diseases [34, 36, 63, 67]. Since opening of KKH in 1982 [4], markets for local products started to develop, thus exerting pressure on natural resources [33, 68].

Ethnobotanical survey

The study was carried out from January 2017 to May 2018, during which 300+ local community members



were approached via focused group discussions (FGDs), semi-structured interviews, and homework assignments for 8th to 12th Grade school and college students. The participants included men and women community members, local traditional health practitioners (THPs), community elders, and wholesalers and retailers of medicinal plants. Following the snowball sampling methodology [69–74], a total of 15 FGDs and 240 individual interviews were conducted for data collection. Sixty students were provided with a set of questionnaires to reach their families and understand what kind of plants/herbal remedies are used in the families and how traditional knowledge is preserved within a family. This method was tested in Nepal and was quite effective to document traditional knowledge [75]. Prior permission and consent for data collection and publication was obtained from all the participants. The homework assignments for school students were administered under the supervision of concerned class teachers. Data on age and gender was also acquired from the participants. These methods resulted in data covering an ethnobotanical inventory of plants, part used, therapeutic utilities, the location and timing of acquiring the resource, and the existing markets with trade opportunities. The initial taxonomic identification of medicinal plants was done by a taxonomist in the field [76–78] and by cross-referencing photographs, voucher specimens, and the local name of species with previously available material and literature from the study area [9, 35, 41, 79, 80]. Information on local names of plant species, and parts of the plant utilized for different medicinal purposes was recorded. Collected specimens were mounted on herbarium sheets and identified by the taxonomists in the Karakoram International University. The voucher specimens were authenticated through the plant list (www.theplantlist.org), tropicos (www.tropicos.org), and flora of Pakistan (www.eflora-s.org) and deposited in the herbarium of the Karakoram International University.

Quantitative analysis

We regressed age of participants against the number of species reported by respective age group and people engaged in the trade of medicinal plants during the survey to understand TK in the region. We used CoKriging method [81] to interpolate the number of species recorded and used in various medical conditions from the surveyed locations to recognize section of highest medicinal plant use in the region. Kriging is an advanced geostatistical procedure that generates an estimated surface from a scattered set of points with z values. CoKriging is multivariate extension of Kriging method that uses information from one or more correlated variables measured in the same range. This method is useful in mountainous areas with rugged terrain. This

method has been tested and used in tree richness mapping, abundance mapping, and recognizing areas at high risk of species invasion [81–83].

We calculated quantitative measures like informant consensus factor (ICF), use value (UV), and relative frequency of citation (RFC) for medicinal plants based on 208 illnesses categorized into 29 ailment groups from each district in the survey. The illnesses were categorized based on a particular part of the body affected or particular illnesses for multiple parts of the body. For example, all kinds of skin illnesses were categorized under skin infections while diseases related to stomach and intestinal disorders were grouped into stomach and intestine category (further detail is available in Additional files 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, and 12). A brief description of the quantitative measures is provided below:

Informant consensus factor (ICF)

The ICF was calculated to find a consensus between participants on the reported treatments for each set of diseases [79, 84, 85]. ICF was calculated using given formula [79, 86, 87];

$$ICF = \frac{(Nur-Nt)}{(Nur-1)}$$

where Nur indicates the number of use reports for a specific disease category and Nt mentions the number of taxa used for the disease category.

Use value (UV)

Use value determines the quantitative measure for relatively important local plant species [78, 87–89]. The use value is calculated using the following formula:

$$UV = \frac{\sum U_i}{N}$$

where $\sum U_i$ is the total number of use reports for a given species and N is the total number of participants inquired for the species.

Relative frequency of citation (RFC)

Ethno medicinal data was quantitatively analyzed using RFC which indicated the local importance of medicinal species. The RFC was calculated using given formula [87, 89, 90]:

$$RFC = \frac{FC}{N} (0 < RFC < 1)$$

where FC is the number of participants reporting on the use of a plant species and N is the total number of participants in the survey.

Discriminant analysis (DA)

We used discriminant analysis to delimit the geographical differences of the ethnobotanical knowledge using quantitative measures (viz. UV and RCF), highly used species, and number of treatments. Discriminant function analysis or discriminate analysis (DA) determines naturally occurring groups and the variables responsible for segregating amongst them [91–93].

Results and discussion

Demographic features of the participants

More than 300 participants including 70% men and 30% women were interviewed during the survey (Fig. 2). The Yashkun, Sheen/Shinaki, Burushos, Wakhi, and Balti communities approached during the survey possessed good knowledge on medicinal plants use. Although these tribes maintain individual identities representing different parts of the mountainous region, the cross-cultural interactions have led to the growth and diversification of the traditional knowledge. The results from FGDs, HH, and market surveys revealed that regardless of ethnicity, THPs and retailers retain a significant level of information on the medicinal plants of the region, the locations from where the plants can be acquired and the ailments they can be utilized for. Although the participants from the community and THPs provided more extensive information than the retailers, when it came to the question of how to use a certain plant as medicine, the retailers had ample information to share on a higher number of medicinal plants and their general uses (Fig. 3 Fig. 4a). We found that participants above 50 years of age had significant traditional knowledge regarding utilization of medicinal plants. This was evident from the number of species and their uses reported per interview where a higher number of species was reported by

participants above 50 years of age. Our results from the linear regression ($R^2 = 0.65$; $p < 0.0001$) also revealed that number of species reported increase with age of the participant (Fig. 4b). The students responding to homework assignments mostly brought information from women representatives of the household, thus providing relevant information on which plant species are kept at home and used for treating common sicknesses.

Taxonomic diversity

We documented a total of 231 species representing 141 genera and 61 families (Table 2). *Compositae* was the most dominant family with 30 (12.9%) species reported, followed by Leguminosae, Lamiaceae, Rosaceae, Polygonaceae, Ranunculaceae, Salicaceae, Apiaceae, and Berberidaceae, with 16 (6.9%), 15 (6.5%), 15 (6.5%), 14 (6.1), 10 (4.3%), 10 (4.3%), 9 (3.9%), and 9 (3.9%) species, respectively. In total, 208 diseases were treated with documented species. The herbaceous flora accounts for 64% of the total reported species followed by shrubs at 20%, trees at 13%, and grasses at 3% (Fig. 5). Figure 6 represents the percentages of uses of different parts of the medicinal plants (further detail is available in Additional files 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, and 12 where a comprehensive list of species is provided besides the district level analysis).

During the survey, the participants revealed that medicinal plants are acquired from special locations and at a specific time, i.e., during a FGD in Upper Hunza (Gujal), the participants mentioned that medicinal plants are found both at the village and in pastures, yet the plants from pastures are used for medicinal purposes as those found in villages are not considered effective for medication. This needs to be scientifically verified from the field




Fig. 2 a Comparison of medicinal plant species highest occurrence and uses in Gilgit-Baltistan region. **b** Linear regression showing relation of age and number of plant species reported. Legend TK = participants and Trade = retailers and traditional healers




Fig. 3 Section of the highest use and occurrence of medicinal plants and uses in Gilgit-Baltistan region

area, yet cases of same plant species presenting different chemical traits has been a known fact [94, 95].

Occurrence and markets for medicinal plants in Gilgit-Baltistan

Interpolated results revealed that the utilization as well as high occurrence of medicinal plants is mainly concentrated in two locations (Fig. 7). Valleys from Skardu, Gilgit, and Ghizer are reported for the highest number of medicinal plants and their uses. There are no formal markets for medicinal plants in the region [35]. Gilgit and Skardu being the main business hubs serve as main markets for wholesale, retail, and purchase of medicinal plants. Besides, small amounts of the medicinal plants are supplied to bigger markets in Rawalpindi and Lahore by the wholesalers. The market players reported Skardu, Astore, and Gilgit districts as the main suppliers of medicinal plants as well.

Participants from the market revealed that most of the large herbal medicine production companies in Pakistan rely on raw materials from Indian Territory representing

the same region across the border. Although a clear percentage of product flow was not known, during the market survey, the participants mentioned that the current markets for locally available plants are limited to the small town market centers and partly target markets in big cities like Rawalpindi (10%) and Lahore (10%) with meager shares moving out across the border to China (1–2%). The participants emphasized the involvement of government agencies for sustainable promotion, collection, and utilization of diverse plant resources in the region.

According to the retailers in Gilgit-Baltistan, *Thymus linearis*, *Delphinium brunonianum*, *Bergenia stracheyi*, *Saussurea heteromalla*, *Saussurea lappa*, *Carthamus tinctorius*, *Peganum harmala*, *Rheum emodi*, *Mentha longifolia*, *Mentha arvensis*, *Valeriana wallichii*, *Berberis lyceum*, and *Elaeagnus rhamnooides* account for the most demanded and utilized species. Although most of the customers were aware of the uses, the retailers had significant knowledge of the plant species they sold. Our survey results revealed that there is a large group of locals in Gilgit-Baltistan who rely on






Fig. 4 Grouping survey sites in Gilgit-Baltistan based on the medicinal plants and their use reports



traditional medicine, yet these practices and the knowledge associated with them is rapidly depleting from the region. All the participants included in the survey had used medicinal plants for treating a medical condition at least once in their life. 79.5% of participants reported a transfer of knowledge from family elders. Out of these, 48% were above 50 years, 38% above 40 years while only 13% were below 40 years of age. 29.1% reported learning from community elders, of whom 64% were above 50 years, 32% above 40 years and only 3% below 40 years. 15.6% reported that they had acquired knowledge through interaction with other people, including THPs, plant collectors, and traders from other communities that include 87% above 50 years and 13% above 40 years. 20.1% were not aware of any treatments through medicinal plants which included only 4 and 26% participants above 50 and 40 years of age respectively while 70% of these participants fell below 40 years of age. Our results clearly indicate that the main source of knowledge transfer rests within the family. Yet only 13% of participants below the age of 40 confirmed that they had received such knowledge from within the family. Traditional healers in the community are experts in recognizing and collecting medicinal plants while other community members are not fully aware of the exact timing for collection. This is one of the main reasons why knowledge transfer is mainly through family elders as family secrets are not shared with outsiders. This






Fig. 7 Habitat/life form of species reported during the survey

allows the family to practice special medicinal remedies as well as retain a good image as regular suppliers to the wholesalers, retailers, and THPs in the region. The gradual expansion of trade and an increasing demand for medicinal plants in and outside the region has a positive impact on knowledge sharing. Most of the suppliers involved in the supply chain of medicinal plants represent the age group 40–60 which raises great concern for the future. Those representing the younger age group are either not collecting proper plants at the proper time or are too keen on gaining more financial benefits, thus not taking long-term conservation into account while collecting. This ever increasing gap between the young and old generation is affecting the knowledge transfer mechanism, in combination with over-exploitation and lack of conservation strategies, and the impacts of climatic changes. Such issues have also been observed in other studies [9, 30, 34, 42].

Informant consensus factor (ICF)

The ICF analysis was done separately for each location in order to assess a clearer picture on which diseases stand out at each location and how many plant species are utilized for their treatment. Table 1 includes the ICF values for diseases divided into 29 disease categories from each of the study locations. The table indicates that stomach and intestinal disorders, respiratory disorders, skin infections, internal and external wounds, pain relief, ear, nose and throat disorders, hepatitis, and livestock diseases were the top disease categories reported from all the field sites during the field study. Most of the ailment categories were reported with a high ICF

value based on the number of species and their usage reports. This is an indication of homogeneity in responses of the participants from each of the study sites in terms of the medicinal plant species and the modes of their use for addressing a particular disease. The results from ICF values clearly indicate that diseases related to stomach and intestinal disorders; respiratory disorders especially asthma; skin infections; and ear, nose, and throat infections were most common diseases, which is also supported by various publications [13, 18, 41]. Besides, the communities generally rely on medicinal plants for treating different kinds of internal and external wounds [10, 13, 18, 39, 41]. It is also evident that community members owning livestock hugely depend on traditional mode of medication [38].

Relative frequency of citation (RFC) and use value (UV)

The RFC and UV values for each plant species were calculated in order to validate the frequency of citation for the species used for different ailment categories. The values were calculated at district level in order to authenticate the local frequency of use. The RFC value is used for verifying the use of a medicinal plant species for different diseases while the UV value is an indication for the relative importance of these species in a particular population [78]. The highest RFC value from all the sites was calculated for *Dracocephalum nuristanicum* Rech.f. & Edelb. (0.7) followed by *Cupressus sempervirens* L., *Prunella vulgaris* L., and *Potentilla argyrophylla* Wall. ex Lehm., averaged at 0.47 each indicating that these species were highly reported by the participants of the study. RFC directly depends on the

Table 1 ICF, number of uses, and species used for each disease category

Category of diseases	Central Hunza				Ghizer				Gojal Hunza				Jalalabad							
	ICF	Number of use reports	% age of use report	Number of species used	ICF	Number of use reports	% age of use report	Number of species used	ICF	Number of use reports	% age of use report	Number of species used	ICF	Number of use reports	% age of use report	Number of species used				
Wounds	0.92	103	4.82	9	5.77	0.93	101	4.12	8	3.85	0.89	48	6.16	5.77	0.94	97	3.97	7	4.24	
Skin infections	0.92	163	7.62	14	8.97	0.93	230	9.38	16	7.69	0.89	36	4.62	5	4.81	0.95	256	10.49	15	9.09
Pain relief	0.95	340	15.90	17	10.90	0.92	227	9.25	18	8.65	0.92	49	6.29	5	4.81	0.94	290	11.88	17	10.30
Kidney and uterus	0.95	101	4.72	6	3.85	0.90	99	4.04	11	5.29	0.94	19	2.44	2	1.92	0.96	122	5.00	6	3.64
Diaphoretic	0.00	0	0.00	0	0.00	1.00	7	0.29	1	0.48	0.00	0	0.00	0	0.00	0.00	0	0.00	0	0.00
Others	0.95	82	3.83	5	3.21	0.93	141	5.75	11	5.29	0.88	70	8.99	9	8.65	0.91	36	1.47	4	2.42
Cardiac stimulant	1.00	11	0.51	1	0.64	0.91	36	1.47	4	1.92	0.85	14	1.80	3	2.88	0.94	91	3.73	6	3.64
Stomach and intestine	0.94	305	14.26	18	11.54	0.93	429	17.49	30	14.42	0.90	126	16.17	14	13.46	0.96	236	9.67	11	6.67
Asthma/breathing/respiratory/pulmonary	0.93	180	8.42	14	8.97	0.94	261	10.64	17	8.17	0.90	80	10.27	9	8.65	0.94	134	5.49	9	5.45
Anti-inflammatory	0.92	14	0.65	2	1.28	0.90	62	2.53	7	3.37	0.92	14	1.80	2	1.92	0.91	65	2.66	7	4.24
Cancers	1.00	8	0.37	1	0.64	0.93	44	1.79	4	1.92	0.00	0	0.00	0	0.00	1.00	67	2.74	1	0.61
Cytoprotective	0.92	50	2.34	5	3.21	0.94	124	5.06	8	3.85	0.88	9	1.16	2	1.92	0.94	122	5.00	8	4.85
HIV	0.00	0	0.00	0	0.00	0.00	0	0.00	0	0.00	0.00	0	0.00	0	0.00	0.00	0	0.00	0	0.00
Hepatitis	0.90	94	4.39	10	6.41	0.90	50	2.04	6	2.88	0.85	34	4.36	6	5.77	0.92	66	2.70	6	3.64
Anti-stress/hypertension	0.93	70	3.27	6	3.85	0.93	42	1.71	4	1.92	0.85	14	1.80	3	2.88	0.95	39	1.60	3	1.82
Hepatoprotective/liver	0.94	19	0.89	2	1.28	0.92	27	1.10	3	1.44	0.00	0	0.00	0	0.00	0.91	83	3.40	8	4.85
Ear, nose, and throat	0.94	211	9.86	14	8.97	0.93	177	7.22	14	6.73	0.90	101	12.97	12	11.54	0.94	266	10.90	17	10.30
Menses/diseases	0.90	43	2.01	5	3.21	0.88	17	0.69	3	1.44	1.00	5	0.64	1	0.96	0.93	31	1.27	3	1.82
Brain and nervous disorders	1.00	10	0.47	1	0.64	0.89	28	1.14	4	1.92	0.00	0	0.00	0	0.00	0.93	128	5.24	10	6.06
Weight loss and fat reduction	0.96	26	1.22	2	1.28	0.86	23	0.94	4	1.92	0.90	11	1.41	2	1.92	0.00	0	0.00	0	0.00
Eye diseases	0.96	26	1.22	2	1.28	0.93	15	0.61	2	0.96	0.88	17	2.18	3	2.88	0.93	28	1.15	3	1.82
Diabetes	0.97	32	1.50	2	1.28	0.90	62	2.53	7	3.37	1.00	5	0.64	1	0.96	0.92	39	1.60	4	2.42
Teeth and gums	0.92	39	1.82	4	2.56	0.91	35	1.43	4	1.92	0.00	0	0.00	0	0.00	0.93	45	1.84	4	2.42
Blood purifier/	0.93	84	3.93	7	4.49	0.91	131	5.34	13	6.25	0.91	23	2.95	3	2.88	0.95	111	4.55	7	4.24

Table 1 ICF, number of uses, and species used for each disease category (*Continued*)

Table 1 ICF, number of uses, and species used for each disease category (Continued)

Category of diseases	Kargha				Nagar				Skardu				Astore							
	ICF	Number of use reports	% age of use report	Number of species used	ICF	Number of use reports	% age of use report	Number of species used	ICF	Number of use reports	% age of use report	Number of species used	ICF	Number of use reports	% age of use report	Number of species used				
Wounds	0.89	185	9.36	21	8.71	0.91	139	7.18	13	7.39	0.87	209	9.04	28	9.27	0.89	19	3.85	3	4.23
Skin infections	0.91	138	6.98	13	5.39	0.90	148	7.64	15	8.52	0.90	222	9.60	24	7.95	0.89	45	9.13	6	8.45
Pain relief	0.90	199	10.07	20	8.30	0.93	214	11.05	15	8.52	0.89	203	8.78	23	7.62	0.91	23	4.67	3	4.23
Kidney and uterus	0.86	110	5.56	16	6.64	0.95	118	6.10	7	3.98	0.85	137	5.92	21	6.95	0.89	38	7.71	5	7.04
Diaphoretic	1.00	3	0.15	1	0.41	0.00	0	0.00	0	0.00	0.00	0	0.00	0	0.00	0.00	0	0.00	0	0.00
Others	0.89	20	1.01	3	1.24	0.93	110	5.68	10	5.68	0.00	0	0.00	0	0.00	0.00	0	0.00	0	0.00
Cardiac stimulant	0.85	14	0.71	3	1.24	0.95	20	1.03	2	1.14	0.89	10	0.43	2	0.66	0.00	0	0.00	0	0.00
Stomach and intestine	0.90	483	24.43	51	21.16	0.91	271	14.00	24	13.64	0.90	638	27.58	63	20.86	0.87	107	21.70	15	21.13
Asthma/breathing/respiratory/pulmonary	0.90	145	7.33	15	6.22	0.93	122	6.30	9	5.11	0.86	210	9.08	30	9.93	0.90	68	13.79	8	11.27
Anti-inflammatory	0.84	51	2.58	9	3.73	0.90	41	2.12	5	2.84	0.83	42	1.82	8	2.65	0.00	0	0.00	0	0.00
Cancers	0.93	41	2.07	4	1.66	0.00	0	0.00	0	0.00	0.00	0	0.00	0	0.00	1.00	5	1.01	1	1.41
Cytoprotective	0.92	66	3.34	6	2.49	0.92	40	2.07	5	2.84	0.89	20	0.86	3	0.99	1.00	5	1.01	1	1.41
HIV	0.00	1	0.05	1	0.41	0.00	0	0.00	0	0.00	0.00	0	0.00	0	0.00	0.00	0	0.00	0	0.00
Hepatitis	0.78	38	1.92	9	3.73	0.92	65	3.36	6	3.41	0.85	49	2.12	8	2.65	0.85	41	8.32	7	9.86
Anti-stress/hypertension	0.87	16	0.81	3	1.24	0.92	40	2.07	4	2.27	0.88	33	1.43	5	1.66	1.00	14	2.84	1	1.41
Hepatoprotective/liver	0.86	23	1.16	4	1.66	0.96	24	1.24	2	1.14	0.81	28	1.21	6	1.99	1.00	4	0.81	1	1.41
Ear, nose, and throat	0.87	143	7.23	19	7.88	0.95	169	8.73	10	5.68	0.84	139	6.01	24	7.95	0.86	43	8.72	7	9.86
Menses/diseases	0.90	11	0.56	2	0.83	0.90	50	2.58	6	3.41	0.90	11	0.48	2	0.66	1.00	5	1.01	1	1.41
Brain and nervous disorders	1.00	5	0.25	1	0.41	0.81	17	0.88	4	2.27	0.86	8	0.35	2	0.66	0.00	0	0.00	0	0.00
Weight loss and fat reduction	1.00	7	0.35	1	0.41	0.91	45	2.32	5	2.84	0.89	20	0.86	3	0.99	0.00	0	0.00	0	0.00
Eye diseases	0.93	15	0.76	2	0.83	0.94	17	0.88	2	1.14	0.83	25	1.08	5	1.66	0.00	0	0.00	0	0.00
Diabetes	0.87	16	0.81	3	1.24	0.90	64	3.31	7	3.98	0.82	29	1.25	6	1.99	0.00	0	0.00	0	0.00
Teeth and gums	0.85	21	1.06	4	1.66	0.92	26	1.34	3	1.70	0.88	41	1.77	6	1.99	0.88	26	5.27	4	5.63
Blood purifier/	0.86	64	3.24	10	4.15	0.93	69	3.56	6	3.41	0.87	125	5.40	17	5.63	0.87	31	6.29	5	7.04

Table 1 ICF, number of uses, and species used for each disease category (Continued)

Category of diseases	Kargha			Nagar			Skardu			Astore										
	ICF	Number of use reports	% age of use report	ICF	Number of use reports	% age of use report	ICF	Number of use reports	% age of use report	ICF	Number of use reports	% age of use report								
diseases																				
Vomiting/nausea/ altitude sickness	0.00	0	0.00	1.00	7	0.36	1	0.57	1.00	7	0.30	1	0.33	1.00	7	1.42	1	1.41		
Livestock diseases	0.89	132	6.68	15	6.22	0.88	75	3.87	10	5.68	0.87	102	4.41	14	4.64	0.91	12	2.43	2	2.82
Sexual diseases/ stimulant	1.00	5	0.25	1	0.41	1.00	4	0.21	1	0.57	0.00	0	0.00	0	0.00	0	0.00	0	0.00	
Hemorrhoids/ piles	0.88	25	1.26	4	1.66	0.94	33	1.70	3	1.70	1.00	5	0.22	1	0.33	0.00	0	0.00	0	0.00
Maternal health	0.00	0	0.00	0	0.00	1.00	8	0.41	1	0.57	0.00	0	0.00	0	0.00	0	0.00	0	0.00	

number of participants mentioning use of a particular plant (FC); therefore, the abovementioned plants were very commonly used in the study area. The UV values for *Amaranthus viridis* L. (0.33), *Artemisia herba-alba* Asso (0.28) and *Astragalus zanskarensis* Bunge (0.25), and *Aconitum violaceum* Jacquem. Ex. Stapf (0.25) were the highest averaged from the field sites. The results indicate the usage and reliance on medicinal plants for treatment of multiple diseases. Such reliance on medicinal plants in both humans and livestock are reported from the region [9, 16, 18, 30, 34, 38, 41]. Table 2 includes a list of reported species from all the six districts of Gilgit-Baltistan with their local names, parts used, mode of their utilization, the average values for RFC and UV, number of ailment categories addressed, and the number of uses reported for each of the plant species. Area wise details for each of these species is provided in Additional files 2, 3, 4, 5, 6, 7, 8, and 9 while the detailed list of diseases categorized in 29 ailment categories is provided in S12.

Medicinal systems and affiliations

This region has already witnessed invasion by different cultures and practices [3, 46–51]. The passes created by the Indus River system in Hunza, Shigar, Shyok, Ghizer, Gilgit, and Astore valleys served as the main travel routes for such invasions and exchanges as Gilgit got its famous name “gate to India” [3]. This region witnessed influence from Chinese, Tibetan, ancient Indian, and Unani systems [3, 46]. This influence was reflected during the field survey while recording the uses and modes of use of medicinal plants.

Fifty percent of the participants were able to answer the question related to influence of medicinal systems on the indigenous knowledge existing in the region; where 28% referred to Chinese influence, 23% chose Indian subcontinent while 18 and 14% selected Scythian/Transoxianan, and ancient Greek influence on the traditional medicinal system. It is worth mentioning that 41% of these participants referred to a mix of at least two of these systems in the current traditional medicinal system. Most of the participants from Astore and Skardu mentioned Indian subcontinent followed by Scythian/Transoxianan and ancient Greek influence while participants from Hunza, Nagar, Gilgit, and Ghizer ranked Chinese influence on top followed by Indian subcontinent, Scythian/Transoxianan, and ancient Greek. The market players on the other hand opted for a mix of all these systems together as they deal with customers from the whole region and are exposed to all medicinal systems prevailing in the region. Another reason for a mixed system described by the market players is the fact that these markets were traditionally placed at regional centers, thus were exposed to THPs

representing different systems. Their experiences with these THPs and fulfilling their demands made them acquire traits from all the systems. Although a clear boundary could not be drawn between these systems, it is likely that medicinal practices in Hunza, Nagar, Ghizer, and Gilgit were influenced by traditional Chinese medicine (TCM) system while the remaining part of study area was dominated by a mix of Ayurveda and Unani systems.

Discriminant analysis (DA)

Discriminant analysis revealed that Astore was distinct from other surveyed areas whereas some overlap can be observed in other surveyed areas (Fig. 8). Gojal showed similarity with Skardu and Kargha, while Jalalabad showed few similarities with Kargha and Hunza. Hunza, Nagar, and Ghizer showed very similar traits while sharing few similarities with Skardu, Kargha, and Jalalabad. This analysis mainly considered top 10 medicinal plants used and the parameters calculated from each location. This does not represent linkage with healing systems but separates geographical location, tribal representation, and connectivity to other parts of the region, e.g., China, India, and routes to Western Asia. It also reflects on migration and integration of the local tribes as well as those coming from outside and provides an insight on the influence of different invaders, travelers, and businessmen.

Plant species have responded to latitudinal and elevational changes in their habitat and adjusted over time, yet the increasingly rapid pace of these changes is challenging their adaptability and ability to respond [96–98]. The association of traditional knowledge with these species, trade potentials, and transfer of knowledge from old to new generation will have a direct effect on the conservation of plant species and associated TK [90, 99–103].

Key discussion points

The environment and climatic conditions of Gilgit-Baltistan make it geographically one of the best locations for growth and nourishment of medicinal plants [6, 9, 34, 104]. Traditional medicinal practices hold a significant place in the lives of the local communities. The markets at Gilgit and Skardu are serving as trade centers for medicinal plants from the whole region, and the wholesalers stationed here are responsible for small-scale trade in important plant species in both local and national markets. The study shows that most of the large herbal medicine production companies in Pakistan rely on supply of medicinal plants from Indian territory—representing the same region across the border—indicating that the production and trade of medicinal plants is well organized across the border. The trade of medicinal plants in Pakistan is informal, with little to none state interventions and incentives [35, 105]. The high marketability of medicinal plants has

Table 2 RFC, UV, number of uses, and ailments of species from each location

S no.	Family	Species	Local name	Part used	Mode of use	Location (Astore-Astore, Ghizer-Ghizer, Gojal-Gojal, Central Hunza-Hunza, Jalalabad-Jalalabad, Kargha-Kargha, Nagar-Nagar, Skardu-Skardu)	Average of RFC	Average of UV	Ailment categories	No. of use responses	Voucher no.	Previous citation
1	Amaranthaceae	<i>Aerva lanata</i> (L.) Juss.	Shutpask	Whole plant	Ash (H, N, K, Gh, A), decoction (S, A, Gh, H, K)	Gojal	0.175	0.214286	3	14	MAS-089	17
2	Achyranthes aspera L.	Iaghakay	Flower	Paste	Kargha	0.2	0.176471	3	17	MAS-238	40, 41	
3	Chenopodium album L.	Snew, Sheleet Kunaw	Leaf (S), whole plant (H, Gh)	Paste, infusion, poultice, decoction	Ghizer, Gojal, Hunza, Skardu	0.21875	0.136706	8	59	MAS-121, 9, 12, MAS-153, 13, 17, MAS-457 33, 37		
4	<i>Dysphania botrys</i> (L.) Mosyakin & Clements	Khamra, Khord	Aerial (K), whole plant (S, H)	Powder, decoction	Gojal, Jalalabad, Skardu	0.194444	0.12619	7	57	MAS-128, 35, MAS-229, MAS-482		
5	<i>Amaranthus viridis</i> L.	Dhindo	Leaf	Paste, direct	Kargha, Nagar	0.1	0.333333	3	10	MAS-314, 14, MAS-376		
6	<i>Allium hurtile</i> Kunth	Chung	Bulb	Infusion, direct	Skardu	0.266667	0.176471	3	17	MAS-381, 12, 13, MAS-346, MAS-424		
7	<i>Allium carolinianum</i> DC.	Kachpauk Booma, Chong	Bulb (K, S, A), leaf, bulb (H)	Paste (S), direct, decoction	Astore, Gojal, Kargha, Nagar, Skardu	0.205	0.22988	21	102	MAS-012, 12, 13, MAS-112, 17, MAS-346, MAS-424		
8	<i>Allium cepa</i> L.	Ghashoo, Xong, Song	Bulb (K), leaf, bulb (S), poultice, bulb (H)	Poultice, decoction, direct (H, S, Gh, N, K), paste (S, A, K, Gh)	Hunza, Kargha, Nagar, Skardu	0.16875	0.202381	9	50	MAS-179, 9, 33, MAS-305, 38, 39, MAS-365, MAS-489		
9	<i>Allium sativum</i> L.	Zgoqpa, Bukpa	Bulb	Direct	Hunza, Kargha, Nagar, Skardu	0.31875	0.110367	11	149	MAS-180, 9, 33, 39, MAS-306, MAS-366, MAS-490		
10	Anacardiaceae	<i>Pistacia mutica</i> Fisch. & C.A.Mey.	Daraaw	Branches	Oil	Hunza	0.2	0.142857	1	7	MAS-137	9, 33
11		<i>Pistacia khinjuk</i> stocks	Kakavomn	Galls, resin, wood, leaf	Direct, decoction	Jalalabad	0.175	0.071429	6	84	MAS-203	15
12	Apiaceae	<i>Heracleum candicans</i> Wall. ex DC.	Ghang	Leaf	Decoction	Skardu	0.266667	0.071429	1	14	MAS-392	12
13		<i>Pimpinella diversifolia</i> DC.	Kohniod	Whole plant	Powder, decoction	Astore, Kargha, Skardu	0.269444	0.15	9	60	MAS-028, 13, MAS-270, MAS-440	
14		<i>Pleurospermum candollei</i> (DC) C.B. Clarke in Hook f.	Braqshundun	Whole plant	Decoction	Astore, Skardu	0.35	0.142857	2	14	MAS-037, 13, MAS-449	
15		<i>Angelica glauca</i> Edgew	Choro, Chora	Root (K), stem, seed, root (Gh)	Decoction, powder, direct	Ghizer, Kargha	0.216667	0.166667	10	55	MAS-078, 11, 16, MAS-284	
16		<i>Carum carvi</i> L.	Fili佐oh, Zera, Hayyo	Seed (K, S), seed, fruit (Gh)	Decoction, powder, direct (Gh)	Ghizer, Kargha, Skardu	0.205556	0.139184	8	66	MAS-084, 11, 12, MAS-290, 16	

Table 2 RFC, UV, number of uses, and ailments of species from each location (Continued)

S no.	Family	Species	Local name	Part used (Kargha-K, Nagar-N, Skardu-S, Ghizer-Gh, Hunza-H, Astore-A, Gilgit-G)	Mode of use (Kargha-K, Nagar-N, Skardu-S, Ghizer-Gh, Hunza-H, Astore-A, Gilgit-G)	Location (Astore-Astore, Ghizer-Ghizer, Gojal-Gojal, Central Hunza-Hunza, Jalalabad-Jalalabad, Kargha-Kargha, Nagar-Nagar, Skardu-Skardu)	Average of RFC	Average of UV	Ailment categories	No. of use responses	Voucher no.	Previous citation
17		<i>Daucus carota</i> L.	Phopuce, Jangli, Jut Ghahsoon, Ghasoona, Ghofatwi lona	Leaf, seed (H, N), leaf (S)	Direct, decoction	Hunza, Kargha, Nagar, Skardu	0.291667	0.134921	7	57	MAS-181, 9, 32, 33	
18		<i>Coriandrum sativum</i> L.	Ausu, Naski	Seed	Decoction, direct	Hunza, Kargha, Skardu	0.213889	0.11929	4	41	MAS-307, MAS-491	
19		<i>Foeniculum vulgare</i> Mill.	Badian	Fruit (K), seed (S)	Decoction	Kargha, Skardu	0.2375	0.099206	2	25	MAS-326, 38, 39	
20		<i>Heracleum sphondylium</i> C.B. Clarke	Hiltreet	Leaf	Direct	Kargha, Skardu	0.295833	0.142857	2	14	MAS-328, 39	
21	Asteraceae	<i>Allardia tonentosa</i> Decne.	Talkham	Leaf, flower	Grinded	Skardu	0.2	0.230769	3	13	MAS-380, 39	
22		<i>Leontopodium leontopodium</i>	Naqoshoto	Seed	Decoction	Kargha, Skardu	0.341667	0.098086	4	41	MAS-329, 39	
23	Berberidaceae	<i>Podophyllum emodi</i> Wall. ex Hook.f. & Thomson	Shingoy	Root, rhizome	Decoction	Kargha	0.2	0.2	2	10	MAS-245, 11, 53	
24		<i>Berberis pseudumbellata</i> R.Parker	Ishkeen, Shokurum, Skyrboo	Root, stem, bark (K), whole plant (S), flower, fruit, seed (A)	Decoction, powder	Astore, Jalalabad, Kargha, Skardu	0.325	0.114213	10	111	MAS-019, 13, 15, MAS-220, 19, 27, MAS-261, 34, 39	
25		<i>Berberis lychnum</i> Royle	Zolg, Ishkeen, Skyrboo	Root, leaf seed, bark, fruit, flower (S, K, N), root, leaf, fruit (H), root, leaf, stem, fruit (Gh)	Decoction	Ghizer, Gojal, Hunza, Kargha, Nagar, Skardu	0.322222	0.092532	55	743	MAS-061, 9, 14, MAS-118, 16, 17, MAS-351, 32, 33, MAS-454, 34, 37, 38	
26		<i>Berberis brandisiana</i> Ahrendt	Ishkeen, Ishkenachi	Root, stem, bark	Decoction	Jalalabad, Kargha, Nagar, Skardu	0.33125	0.114998	39	490	MAS-235, 15, 34	
27		<i>Berberis orthotropis</i> Bien. ex Aitch.	Ishkeen, Skyurboo	Root, stem, bark (K), root, stem (S)	Decoction	Jalalabad, Kargha, Skardu	0.341667	0.112444	15	121	MAS-236, 15, 34, MAS-313, MAS-498	
28		<i>Berberis parkeriana</i> C.K.Schneid.	Ishkeen, Skyurboo	Root, stem	Decoction	Kargha, Skardu	0.354167	0.154762	5	33	MAS-317, 34	
29		<i>Berberis stewartiana</i> Jafri	Ishkeen, Shokurum	Root, leaf	Decoction	Kargha, Skardu	0.25	0.224599	6	28	MAS-318, 34	
30		<i>Berberis ulicina</i> Hook.f. & Thomson	Ishkeen, Shokurum	Root, leaf	Decoction	Kargha, Skardu	0.2625	0.204861	5	25	MAS-319, 34	
31		<i>Berberis vulgaris</i> L.	Ishkeen, Shokurum	Fruit (S), leaf, fruit (K)	Decoction, direct	Kargha, Skardu	0.304167	0.154762	6	39	MAS-320, 12, 34, MAS-505, 39	

Table 2 RFC, UV, number of uses, and ailments of species from each location (Continued)

S no.	Family	Species	Local name	Part used (Kargha-K, Nagar-N, Skardu-S, Ghizer-Gh, Hunza-H, Astore-A, Gilgit-G)	Mode of use (Kargha-K, Nagar-N, Skardu-S, Ghizer-Gh, Hunza-H, Astore-A, Gilgit-G)	Location (Astore-Astore, Ghizer-Ghizer, Gojal-Gojal, Central Hunza-Hunza, Jalalabad-Jalalabad, Kargha-Kargha, Nagar-Nagar, Skardu-Skardu)	Average of RFC	Average of UV	Ailment categories	No. of use responses	Voucher no.	Previous citation
32	Betulaceae	<i>Betula utilis</i> D.Don	Xuxi Halli, Jowzee, Furze, Stipa	Bark, wood (N, K, H), bark (S)	Decoction, direct	Gojal, Hunza, Jalalabad, Kargha, Nagar, Skardu	0.2375	0.099554	13	143	MAS-123, 9, 11, MAS-167, 13, 15, MAS-225, 17, 39, MAS-477 53	
33	Boraginaceae	<i>Onosma hispida</i> Wall. ex G. Don	Kangmar	Whole plant	Decoction	Astore, Kargha	0.291667	0.162338	5	32	MAS-020, 11, 13, MAS-262	
34	Brassicaceae	<i>Brassica oleracea</i> var. <i>botrytis</i> L.	Phul Gobi	Flower	Direct	Hunza	0.175	0.166667	1	6	MAS-132 9	
35		<i>Descurainia sophia</i> (L.) Webb ex Plant	Khashir	Whole plant	Powder, decoction	Astore, Skardu	0.233333	0.174242	4	23	MAS-034, 13, MAS-446	
36		<i>Raphanus sativus</i> L.	Moolo, Gholaifi	Leaf sonna	Direct	Hunza, Kargha, Nagar, Skardu	0.2875	0.071584	5	99	MAS-182, 9, 39, MAS-308, MAS-368, MAS-492	
37		<i>Brassica oleracea</i> var. <i>capitata</i> L.	Band Gobi	Flower	Direct	Hunza, Nagar	0.15	0.133333	2	16	MAS-186, 9, MAS-369	
38		<i>Brassica juncea</i> (L.) Czern.	Sarsung mar	Seed	Oil	Kargha, Skardu	0.191667	0.142857	2	14	MAS-323, 39, MAS-508	
39		<i>Lepidium latifolium</i> L.	Sonna	Leaf (K), leaf, root (S)	Powder, infusion (S)	Kargha, Skardu	0.3375	0.171429	4	22	MAS-330, 12, MAS-515	
40	Buxaceae	<i>Buxus papyrifera</i> C.K. Schneid	Angaroo	Leaf	Oil	Skardu	0.233333	0.083333	1	12	MAS-388 38	
41	Campanulaceae	<i>Codonopsis clematidea</i> (Schrenk) C.B.Clarke	Loo sunma/Bajoo mindoq	Flower		Astore, Skardu	0.366667	0.107143	2	21	MAS-032, 12, 13, MAS-444	
42	Cannabaceae	<i>Cannabis sativa</i> L.	Thoonch	Seed (N, H), whole plant (K)	Direct	Hunza, Kargha, Nagar	0.258333	0.132762	12	101	MAS-178, 9, 14, MAS-304, 33, 53, MAS-364	
43	Capparaceae	<i>Capparis spinosa</i> L.	Kraba, Kavir, Kappar, Chopir, Short, Champaarang, Thoonch	Root, bark fruit, seed, branches, flower (N, K), root, fruit seed (S), seed, flower, fruit (H)	Oil (H, N, K), powder, decoction, oil (S), paste (H)	Gojal, Hunza, Jalalabad, Kargha, Nagar, Skardu	0.276389	0.093599	25	302	MAS-124, 9, 12, MAS-226, 15, 17, MAS-339, 33, 37, MAS-478 53	
44	Caprifoliaceae	<i>Lonicera microphylla</i> Wild. ex Schult.	Pushkar	Stem, braches, fruit	Paste	Jalalabad	0.2	0.043478	1	23	MAS-199 15	
45		<i>Valeriana wallachii</i> DC.	Mushk-bala	Root (K, H), rhizomes (S)	Powder, decoction, paste	Gojal, Kargha, Skardu	0.266667	0.149335	6	40	MAS-130, 11, 17, MAS-300, 38, MAS-484	
46		<i>Lonicera caprifolium</i> Hook f. & Thomson	Kraba	Leaf	Direct	Kargha, Skardu	0.270833	0.142857	2	14	MAS-331, 39, MAS-516	
47	Caryophyllaceae	<i>Cerastium glomeratum</i>	Bloghar	Whole plant	Direct	Astore, Kargha, Skardu	0.222222	0.118276	6	54	MAS-024, 13	

Table 2 RFC, UV, number of uses, and ailments of species from each location (Continued)

S no.	Family	Species	Local name	Part used (Kargha-K, Nagar-N, Skardu-S, Ghizer-Gh, Hunza-H, Astore-A, Gilgit-G)	Mode of use (Kargha-K, Nagar-N, Skardu-S, Ghizer, Gojal-Gojal, Central Hunza-Hunza, Jalabad-Jalabad, Kargha-Kargha, Nagar-Nagar, Skardu-Skardu)	Average of RFC	Average of UV	Ailment categories	No. of use responses	Voucher no.	Previous citation	
fontanum Baumg.												
48	Compositae	<i>Anaphalis nepalensis</i>	Chikee	Flower, fruit	Dried flower, powder, Ghizer fume (Gh)	0.233333	0.142857	1	7	MAS-039	16	
49		<i>Artemisia annua</i> L.	Xoon	Whole plant	Direct	Ghizer	0.233333	0.157895	3	19	MAS-040	37
50		<i>Artemisia dubia</i> Wall. Ex Bess.	Bursay	Whole plant	Paste, powder	Ghizer	0.2	0.142857	3	21	MAS-041	37
51		<i>Artemisia herba-alba</i> Asso	Kho Bursay	Whole plant	Decoction	Ghizer	0.2	0.272727	3	11	MAS-042	37
52		<i>Saussurea heteromalla</i> (D.Don) Hand.-Mazz.	Kali zira	Weed	Paste, direct	Ghizer	0.1	0.111111	2	18	MAS-049	16
53		<i>Anaphalis triplinervis</i> (Sims) Sims ex C.B.Clarke	Yeepwoosh	Leaf, flower	Poultice, dried leaf and flower	Gojal	0.225	0.166667	4	24	MAS-090	17
54		<i>Tragopogon dubius</i> Scop	Kreel woosh	Flower	Decoction	Gojal	0.15	0.142857	1	7	MAS-104	17
55		<i>Achillea millefolium</i> L.	Yarrow	Flower	Decoction, poultice	Kargha	0.175	0.142857	3	21	MAS-237	40, 41
56		<i>Artemisia laciniata</i> Willd.	Khampa	Leaf	Paste	Kargha	0.3	0.142857	1	7	MAS-239	11, 12
57		<i>Artemisia rutifolia</i> Spreng. Ex Spreng	Kho Bursay	Aerial	Paste	Kargha	0.175	0.142857	1	7	MAS-240	38
58		<i>Artemisia fragrans</i> Willd.	Kho Bursay	Aerial	Powder	Skardu	0.266667	0.166667	1	6	MAS-384	38
59		<i>Artemisia santolinifolia</i> Turcz. Ex Krásch.	Kho Bursay	Leaf, stem	Powder, paste	Skardu	0.266667	0.2	2	10	MAS-385	39
60		<i>Jurinea dolomiaeæ</i> Boiss.	Sathing	Leaf, root	Decoction, poultice	Skardu	0.233333	0.214286	3	14	MAS-393	12
61		<i>Pseudognaphalium luteoalbum</i> (L.) Hilliard & B.L. Burtt	Thrin	Leaf	Decoction	Skardu	0.366667	0.142857	1	7	MAS-404	12
62		<i>Senecio chrysanthemoides</i> DC.	Api mindaq	Leaf, flower, root	Decoction, poultice	Skardu	0.233333	0.157895	3	19	MAS-409	12
63		<i>Tanacetum senecionis</i> (Jacquem. ex Besser) J.Gay ex DC.	Hiltree/Tialo	Flower	Powder, infusion, decoction	Skardu	0.4	0.166667	3	18	MAS-411	38

Table 2 RFC, UV, number of uses, and ailments of species from each location (Continued)

S no.	Family	Species	Local name	Part used	Mode of use	Location (Astore-Astore, Ghizer-Ghizer, Gojal-Gojal, Central Hunza-Hunza, Jalabad-Jalabad, Kargha-Kargha, Nagar-Nagar, Skardu-Skardu)	Average of RFC	Average of UV	Ailment categories	No. of use responses	Voucher no.	Previous citation
64		<i>Taraxacum officinale</i> (L.) Weber ex F.H.Wigg.	Dodduli, Mamo Shiknachi, Ishkanachi, Shartha, Takting, Khosnas	Leaf root (K, Gh), leaf, flower (S), leaf, latex (H), latex (A)	Decoction, powder (K), infusion (S)	Astore, Ghizer, Gojal, Hunza, Kargha, Skardu	0.222222	0.161198	18	136	MAS-003, MAS-051, MAS-415	9, 13, 14, 16, 17, 33, 37, 38, 53
65		<i>Artemisia brevifolia</i> Wall	Rooner, Bursay, Tarotpesk Bustae	Leaf (N, K), leaf, flower (S, H), whole plant (A, Gh)	Poultice, direct	Astore, Ghizer, Gojal, Kargha, Nagar, Skardu	0.216667	0.218759	12	62	MAS-006, MAS-054, MAS-109	13, 17, 35, 37, 38, 39
66		<i>Artemisia maritima</i> L.	Rooner, Zooon Bursay	Leaf bud, flower (N, K Gh), aerial (H), flower (A)	Direct, paste, decoction (Gh)	Astore, Ghizer, Hunza, Kargha, Nagar, Skardu	0.270833	0.112619	18	228	MAS-007, MAS-055, MAS-142, MAS-419	9, 11, 14, 16, 33, 37, 39, 40, 41, 53
67		<i>Artemisia sieversiana</i> Ehrh.	Hampa, Khampa	Leaf (K, A, N), leaf, flower, root (S)	Infusion, decoction, paste	Astore, Kargha, Nagar, Skardu	0.183333	0.20211	6	29	MAS-022, MAS-264, MAS-349, MAS-434	12, 39
68		<i>Cousinia thomsonii</i> C.B.Clarke	Charchu	Flower	Direct	Astore, Skardu	0.266667	0.142857	2	14	MAS-033, MAS-445	13
69		<i>Carthamus tinctorius</i> L.	Pock, Poong	Flower, seed	Decoction, oil	Ghizer, Hunza	0.370833	0.079828	14	175	MAS-066, MAS-154, MAS-459	9, 16, 32, 33, 37
70		<i>Echinops echinatus</i> Roxb.	Jacheer	Whole plant	Decoction, direct	Ghizer, Hunza, Kargha, Nagar	0.260417	0.103175	6	96	MAS-072, MAS-160, MAS-281, MAS-356	9, 14, 16
71		<i>Saussurea lappa</i> (Decne.) Sch.Bip.	Minal	Root (K), stem, root (Gh)	Powder, paste, decoction	Ghizer, Kargha	0.204167	0.139959	9	75	MAS-082, MAS-288	11, 16, 37, 53
72		<i>Artemisia scoparia</i> Waldst. & Kitam.	Khobustae	Leaf, flower (K, S), whole plant (Gh)	Paste, fume (Gh), decoction (S)	Ghizer, Kargha, Skardu	0.202778	0.176667	6	43	MAS-083, MAS-289	13, 37
73		<i>Cichorium intybus</i> L.	Ishkinachi, Caroop, qarali Chicknachi	Whole plant (N, K), root, leaf (S)	Direct, infusion, decoction, decoction (H)	Gojal, Hunza, Kargha, Nagar, Skardu	0.251667	0.148798	12	99	MAS-126, MAS-297, MAS-360	12, 14, 17, 33
74		<i>Tanacetum gracile</i> Hook.f. & Thomson	Cerpho bursay, serfo bursay, Bursay	Leaf	Decoction, powder (S), direct	Hunza, Kargha, Skardu	0.252778	0.155844	5	32	MAS-185, MAS-311, MAS-495	12, 39, 40, 41
75		<i>Artemisia absinthium</i> L.	Zoon	Whole plant	Infusion, paste, powder	Kargha, Nagar	0.175	0.177778	4	24	MAS-315, MAS-377	9, 11, 14
76		<i>Seriphidium brevifolium</i> (Wall. ex DC.) Ling &	Bursay	Leaf (K), flower (S)	Powder, decoction, direct	Kargha, Skardu	0.320833	0.142857	2	14	MAS-336, MAS-521	12, 39

Table 2 RFC, UV, number of uses, and ailments of species from each location (Continued)

S no.	Family	Species	Local name	Part used (Kargha-K, Nagar-N, Skardu-S, Ghizer-Gh, Hunza-H, Astore-A, Gilgit-G)	Mode of use (Kargha-K, Nagar-N, Skardu-S, Ghizer-Gh, Hunza-H, Astore-A, Gilgit-G)	Location (Astore-Astore, Ghizer-Ghizer, Gojal-Gojal, Central Hunza-Hunza, Jalalabad-Jalalabad, Kargha-Kargha, Nagar-Nagar, Skardu-Skardu)	Average of RFC	Average of UV	Ailment categories	No. of use responses	Voucher no.	Previous citation
77		<i>Tanacetum falconeri</i> Hook.f.	Hattyri, Htialo, Pholing, Zoon, Tyalo	Whole plant (K), leaf (S)	Direct, powder	Kargha, Skardu	0.204167	0.190909	4	21	MAS-338, MAS-523	13, 39
78	Convolvulaceae	<i>Convolvulus arvensis</i> L.	Thringthingmo	Whole plant	Decoction, powder	Astore, Kargha, Skardu	0.197222	0.166667	3	18	MAS-023, MAS-265, MAS-435	13
79		<i>Cuscuta reflexa</i> Roxb.	Ghbul thaq (S)	Stem, flower (K), whole plant (S)	Decoction, direct (S)	Kargha, Skardu	0.2375	0.1625	4	26	MAS-325, MAS-510	38, 39
80	Crassulaceae	<i>Rhodiola imbricata</i> Edgew.	Chundol	Root	Powder	Skardu	0.233333	0.222222	4	18	MAS-406	12
81	Cucurbitaceae	<i>Cupressus sempervirens</i> L.	Sato	Fruit, stem	Decoction, direct	Ghizer	0.466667	0.095238	2	21	MAS-044	16, 37
82		<i>Citrullus vulgaris</i> Schrad.	Bowar	Fruit	Decoction	Hunza	0.225	0.045455	1	22	MAS-133	9, 33
83		<i>Cucurbita pepo</i> L.	Hasar	Seed	Oil, direct	Nagar	0.4	0.085714	3	35	MAS-339	9
84	Cupressaceae	<i>Juniperus squamata</i> Buch. turkestanica Kom. M.Bieb.	Cheleh, Chili, Hasnuk, Yarz, Shukpa	Fruit, twigs	Infusion, oil, paste	Ghizer	0.233333	0.132075	7	53	MAS-045	37
85		<i>Juniperus turkestanica</i> Kom. M.Bieb.	Cheleh	Leaf, fruit, wood	Powder, decoction	Jalalabad	0.175	0.054054	2	37	MAS-198	15
86		<i>Juniperus excelsa</i> M.Bieb.	Cheleh, Chili, Hasnuk, Yarz, Shukpa	Fruit, wood, leaf (N, K, H), fruit (S, A)	Ash, powder, decoction, paste (Gh)	Astore, Ghizer, Gojal, Jalalabad, Kargha-Kargha, Nagar, Skardu	0.245238	0.157465	25	194	MAS-004, MAS-052, MAS-107, MAS-416	12, 13, 14, 15, 16, 17, 37
87		<i>Juniperus communis</i> L.	Mithnary, Oshuk	Fruit, wood (K, A, S), fruit, wood, oil (Gh)	Infusion, decoction, paste, powder (A)	Astore, Ghizer, Jalalabad, Skardu	0.272917	0.145114	13	112	MAS-009, MAS-215, MAS-421	13, 15, 37
88	Elaeagnaceae	<i>Elaeagnus rhamnoidea</i> (L.) A.Nelson	Buru, Buroh, Sebaulkhrom, Soq, Rema, Zakh, Chanso, Karsoq	Leaf, fruit, seed, root, wood (K, N), leaf, fruit seed (S), fruit (H), fruit, stem, leaf (Gh), fruit, leaf (A)	Ash, direct, decoction, powder	Astore, Ghizer, Gojal, Hunza, Jalalabad, Kargha, Nagar, Skardu	0.323958	0.106938	43	747	MAS-002, MAS-050, MAS-105, MAS-414	9, 11, 12, 13, 14, 15, 17, 30, 33, 38, 39, 53
89		<i>Elaeagnus angustifolia</i> L.	Shekarkuch, Gindawar, Ghonair, Sisk, Ghundai	Whole plant (N, K), flower, fruit, gum (Gh, H)	Direct, powder, decoction,	Ghizer, Gojal, Hunza, Jalalabad, Kargha, Nagar	0.295833	0.080555	25	331	MAS-059, MAS-116, MAS-350, MAS-452	9, 14, 15, 16, 17, 32, 33, 37
90	Ephedraceae	<i>Ephedra intermedia</i> Schenck & C.A.Mey.	Shaay Soom	Stem, root	Decoction	Jalalabad	0.2	0.061538	4	65	MAS-194	57
91		<i>Ephedra gerardiana</i> Wall. ex Stapf	Soom, Say, Yemoor, Sopat, Sop	Whole plant (K, Gh), aerial (S), leaf, stem (H)	Decoction	Ghizer, Gojal, Hunza, Jalalabad, Kargha, Skardu	0.294444	0.097941	22	239	MAS-060, MAS-275, MAS-453	9, 11, 12, 14, 15, 16,

Table 2 RFC, UV, number of uses, and ailments of species from each location (Continued)

S no.	Family	Species	Local name	Part used (Kargha-K, Nagar-N, Skardu-S, Ghizer-Gh, Hunza-H, Astore-A, Gilgit-G)	Mode of use (Kargha-K, Nagar-N, Skardu-S, Ghizer, Gojal-Gojal, Central Hunza-Hunza, Jalalabad-Jalalabad, Kargha-Kargha, Nagar-Nagar, Skardu-Skardu)	Average of RFC	Average of UV	Ailment categories	No. of use responses	Voucher no.	Previous citation	
92	Equisetaceae	<i>Equisetum arvense</i> L.	Thangshingy harswa, Thangshing stwa	Aerial (S), whole plant (A)	Decoction	Astore, Skardu	0.283333	0.142857	2	14	MAS-035, MAS-447	32, 33, 37, 38, 53
93	Ericaceae	<i>Rhododendron anthropogonium</i> D. Don	Chauiman	Leaf, flower	Infusion, decoction	Skardu	0.3	0.15	3	20	MAS-407	12
94	Gentianaceae	<i>Swertia petiolata</i> D. Don	Brama	Leaf, root	Paste, decoction, powder (A)	Skardu	0.233333	0.210526	4	19	MAS-410	12
95		<i>Gentiana olivieri</i> Griseb.	Tikta	Leaf, flower (K), leaf (S)	Direct	Kargha, Skardu	0.233333	0.162338	3	18	MAS-327, MAS-512	12, 39
96	Geraniaceae	<i>Geranium nepalense</i> sweet	Bamik	Fruit, root	Poultice, decoction, powder	Skardu	0.3	0.235294	4	17	MAS-391	12
97	Grossulariaceae	<i>Ribes himalaense</i> Royle ex Decne.	Murshatooh	Fruit	Powder	Jalalabad	0.175	0.075	3	40	MAS-204	15
98		<i>Ribes orientale</i> Desf.	Ghonashatooh	Root	Powder	Jalalabad	0.2	0.033333	2	60	MAS-205	15
99		<i>Ribes alpestre</i> Wall. ex Decne.	Shumlooh, Skioruru	Root, flower	Powder, direct	Astore, Gojal, Jalalabad, Skardu	0.189583	0.109821	9	109	MAS-011, MAS-111, MAS-217, MAS-423	13, 15, 17
100	Iridaceae	<i>Crocus sativus</i> L.	Zafran	Flower	Powder	Kargha, Skardu	0.266667	0.142857	2	14	MAS-324, MAS-509	39
101	Juglandaceae	<i>Juglans regia</i> L.	Achow, Ashooh	Root, kernel, wood (N), root, (kernel, seed, wood (K), kernel (S, H))	Oil, direct	Hunza, Jalalabad, Kargha, Nagar, Skardu	0.276667	0.098536	11	139	MAS-173, MAS-301, MAS-488	9, 13, 15, 33
102	Lamiaceae	<i>Nepeta floccosa</i> Benth.	Buzanj	Leaf, flower	Decoction, infusion	Gojal	0.2	0.172414	5	29	MAS-098	17
103		<i>Isodon rugosus</i> (Wall. ex Benth.) Codd	Phaypush	Leaf, branches	Powder	Jalalabad	0.2	0.097561	4	41	MAS-197	15
104		<i>Dracocephalum nuristanicum</i> Rech.f. & Edeh.	Shardun	Leaf, flower, seed	Paste, decoction, infusion	Skardu	0.7	0.064935	5	77	MAS-389	12, 38, 39
105		<i>Mentha haplocalyx</i> Briq.	Shoma	Leaf	Direct, paste	Skardu	0.3	0.214286	3	14	MAS-394	39
106		<i>Nepeta leucoldena</i> Benth. ex Hook.f.	Askuta	Whole plant	Powder, decoction	Skardu	0.366667	0.035714	1	28	MAS-395	12, 38
107		<i>Prunella vulgaris</i> L.	Harswa	Leaf	Decoction	Skardu	0.466667	0.153846	4	26	MAS-403	12
108		<i>Mentha royleana</i>	Foling, Gudunj	Leaf (K, A, S), leaf, flower (H)	Powder, paste (H),	Astore, Gojal, Kargha, Skardu	0.291667	0.104499	9	91	MAS-013,	12, 13,

Table 2 RFC, UV, number of uses, and ailments of species from each location (Continued)

S no.	Family	Species	Local name	Part used (Kargha-K, Nagar-N, Skardu-S, Ghizer-Gh, Hunza-H, Astore-A, Gilgit-G)	Mode of use (Kargha-K, Nagar-N, Skardu-S, Ghizer, Gojal-Gojal, Central Hunza-Hunza, Jalalabad-Jalalabad, Kargha-Kargha, Nagar-Nagar, Skardu-Skardu)	Average of RFC	Average of UV	Ailment categories	No. of use responses	Voucher no.	Previous citation
109		<i>Thymus linearis</i> Benth.	Tumuro, Tumburu, Tumburuk	Whole plant (N, S), leaf, flower (H), flower (A)	Decoction, infusion	Astore, Gojal, Nagar, Skardu	0.222917	0.100514	12	151	MAS-014, MAS-114, MAS-347, MAS-426
110		<i>Mentha longifolia</i> (L.) L.	Fileel, Whadan, Phileel	Leaf, flower (H, K, N), leaf (Gh)	Powder, paste, decoction (Gh)	Ghizer, Gojal, Kargha, Nagar	0.252083	0.12514	29	319	MAS-065, MAS-122, MAS-278, MAS-353
111		<i>Mentha syvestris</i> L.	Bundoo	Leaf, flower	Decoction	Ghizer, Hunza	0.241667	0.101515	8	74	MAS-067, MAS-155, MAS-460
112		<i>Thymus serpyllum</i> L.	Tumuro, Ree tumburuk	Whole plant (K, Gh), leaf, flower (S), aerial (H)	Decoction	Ghizer, Hunza, Kargha, Skardu	0.33125	0.079512	17	268	MAS-074, MAS-074, MAS-162, MAS-283, MAS-467
113		<i>Mentha arvensis</i> L.	Peeno	Leaf (S, K), whole plant (Gh)	Powder, direct, paste	Ghizer, Kargha, Skardu	0.275	0.138889	6	44	MAS-085, MAS-291, MAS-473
114		<i>Perovskia abrotanoides</i> Kar.	Fairing bursay	Flower	Infusion	Hunza, Kargha, Skardu	0.202778	0.126246	6	57	MAS-184, MAS-310, MAS-494
115		<i>Mentha spicata</i> L.	Podina	Whole plant	Decoction	Hunza, Nagar	0.3375	0.046801	8	160	MAS-189, MAS-372
116		<i>Stachys tibetica</i> Vatke	Khampa	Leaf	Powder, direct	Kargha, Skardu	0.304167	0.049043	2	41	MAS-337, MAS-522
117	Leguminosae	<i>Astragalus frigidus</i> (L.) A.Gray	Shashal	Leaf, stem	Powder	Astore	0.233333	0.176471	3	17	MAS-001
118		<i>Astragalus falconeri</i> Bunge	Hapocho	Leaf, stem	Powder	Ghizer	0.233333	0.083333	1	12	MAS-043
119		<i>Melilotus officinalis</i> (L.) Pall.	Bissasing	Whole plant	Decoction	Ghizer	0.333333	0.115385	6	52	MAS-046
120		<i>Astragalus strictus</i> Benth.	Zhop/Thope	Leaf, flower	Direct	Gojal	0.175	0.142857	1	7	MAS-091
121		<i>Melilotus alba</i> Ledeb.	Sinjhi	Aerial	Paste	Gojal	0.175	0.166667	2	12	MAS-096
122		<i>Caragana brevifolia</i> Kom.	Hapochoo	Root	Direct, decoction	Jalalabad	0.175	0.058824	1	17	MAS-191
123		<i>Caragana tragacanthoides</i> var. <i>himalaica</i> Komarov	Hapochoo	Root	Direct, decoction	Jalalabad	0.225	0.037037	1	27	MAS-192

Table 2 RFC, UV, number of uses, and ailments of species from each location (Continued)

S no.	Family	Species	Local name	Part used	Mode of use	Location (Astore-Astore, Ghizer-Ghizer, Gojal-Gojal, Central Hunza-Hunza, Jalalabad-Jalalabad, Kargha-Kargha, Nagar-Nagar, Skardu-Skardu)	Average of RFC	Average of UV	Ailment categories	No. of use responses	Voucher no.	Previous citation
124		<i>Robinia pseudoacacia</i> L.	Kekar	Resin, wood, legumes	Paste	Jalalabad	0.35	0.076923	2	26	MAS-206	15
125		<i>Astragalus zanskarensis</i> Bunge	Shukpa	Leaf, stem	Paste, ash	Skardu	0.2	0.25	2	8	MAS-386	39
126		<i>Trifolium fragiferum</i> L.	Gul-e-Naseen	Leaf, flower	Direct	Skardu	0.233333	0.142857	2	14	MAS-413	38
127		<i>Trifolium pratense</i> L.	Chita-batta, Ol, Jangli shaftal	Flower	Powder	Astore, Hunza, Kargha, Skardu	0.2125	0.078571	6	76	MAS-017, MAS-147, MAS-260, MAS-429	9, 13, 14, 40, 41
128		<i>Cicer microphyllum</i> Benth.	Strajungstwa	Whole plant		Astore, Kargha, Skardu	0.230556	0.087446	3	35	MAS-025, MAS-267, MAS-437	13
129		<i>Medicago sativa</i> L.	Ucharq, Ishfit	Whole plant	Direct, powder, decoction	Ghizer, Gojal	0.2	0.134921	9	70	MAS-058, MAS-115	16, 17, 37
130		<i>Sophora mollis</i> (Royle) Baker	Khakhul, Popshing, Pushool	Leaf (K, Gh), leaf, seed (S), whole plant (H)	paste, powder, decoction (S)	Ghizer, Gojal, Hunza, Kargha, Skardu	0.243333	0.10857	10	108	MAS-119, MAS-151, MAS-277, MAS-455	9, 12, 16, 17, 38, 39
131		<i>Glycyrrhiza glabra</i> L.	Shalako	Root, rhizome (Gh), rhizome (K)	Decoction, paste	Ghizer, Kargha	0.3	0.091954	12	145	MAS-079, MAS-285	11, 16, 37
132		<i>Trigonella foenum-graecum</i> L.	Shamilik	Leaf (K), whole plant (S, Gh)	Direct, decoction	Ghizer, Kargha, Skardu	0.277778	0.111683	6	62	MAS-086, MAS-292, MAS-474	37, 38, 39
133		<i>Astragalus psilocentrus</i> Fisch.	Bjowacharchu, Biacharchoo, Sokhnu, Haipocho	Leaf, stem (K, H), leaf, root, thorny branches (S)	Decoction, infusion (S)	Ghizer, Hunza, Kargha, Skardu	0.258333	0.153501	9	63	MAS-088, MAS-166, MAS-294, MAS-476	9, 12, 13, 38
134	Linaceae	<i>Linum usitatissimum</i> L.	Human	Seed	Powder	Hunza	0.175	0.07874	10	127	MAS-134	9, 32, 33
135	Lythraceae	<i>Punica granatum</i> L.	Danooh, Sio, Dolum, Danu	Fruit, root (K, S), flower, fruit, seed, bark (H)	Decoction, paste, powder, direct	Gojal, Hunza, Jalalabad, Skardu	0.26875	0.112524	19	220	MAS-125, MAS-169, MAS-227, MAS-479	9, 15, 17, 33, 38
136	Malvaceae	<i>Malva neglecta</i> Wall.	Shanishah	Whole plant	Powder, decoction	Gojal	0.125	0.071429	1	14	MAS-095	17
137		<i>Abelmoschus esculentus</i> (L.) Moench	Bhindi	Seed, fruit	Infusion	Hunza, Jalalabad, Nagar	0.308333	0.064052	3	57	MAS-174, MAS-233, MAS-362	9
138		<i>Morus nigra</i> L.	Kini Matooch	Whole plant	Decoction, paste, direct	Hunza	0.2225	0.1	3	30	MAS-135	33
139		<i>Ficus carica</i> L.	Faag, Faak	Fruit, stem latex (N, H, K), fruit (Gh)	Poultice, direct, powder, paste	Ghizer, Hunza, Jalalabad, Nagar	0.275	0.096875	16	203	MAS-070, MAS-158, MAS-224,	9, 15, 16, 33

Table 2 RFC, UV, number of uses, and ailments of species from each location (Continued)

S no.	Family	Species	Local name	Part used (Kargha-K, Nagar-N, Skardu-S, Ghizer-Gh, Hunza-H, Astore-A, Gilgit-G)	Mode of use (Kargha-K, Nagar-N, Skardu-S, Ghizer, Gojal-Gojal, Central Hunza-Hunza, Jalalabad-Jalalabad, Kargha-Kargha, Nagar-Nagar, Skardu-Skardu)	Average of RFC	Average of UV	Ailment categories	No. of use responses	Voucher no.	Previous citation
140		<i>Morus alba</i> L.	Marooch, Shae Marooch	Whole plant	Decoction, paste, direct	Hunza, Jalalabad	0.275	0.065763	7	119	MAS-171, 9, 15, MAS-230, 32, 33
141	Nitrariaceae	<i>Peganum harmala</i> L.	Spandur, Isman, Ispandure, Supandour	Whole plant (N, H), seed (Gh, S)	Powder, decoction, paste (H)	Ghizer, Gojal, Hunza, Nagar, Skardu	0.288333	0.097728	14	143	MAS-063, 9, 16, MAS-120, 17, 33, MAS-352, 38, 53, MAS-456
142	Nyctaginaceae	<i>Mirabilis jalapa</i> L.	Gul-e-Abbas	Flower	Paste	Hunza, Kargha	0.1875	0.142857	2	14	MAS-176, 40, 41, MAS-302
143	Oleaceae	<i>Fraxinus hookeri</i> Wenz.	Kasunar	Bark, wood	Decoction	Jalalabad	0.275	0.047619	2	42	MAS-195, 15
144		<i>Fraxinus xanthoxylodes</i> (G.Don) Wall. ex ADC.	Kasunar	Bark, wood	Decoction	Jalalabad	0.175	0.055556	2	36	MAS-196, 15
145		<i>Olea ferruginea</i> Wall. ex Aitch.	Kawoo	Leaf, wood, bark	Direct, decoction	Jalalabad	0.2	0.071429	1	14	MAS-200, 15
146	Onagraceae	<i>Epilobium latifolium</i> L.	Pondol	Leaf, flower	Paste, decoction	Skardu	0.233333	0.071429	1	14	MAS-390, 12
147	Orchidaceae	<i>Dactylochila haragirea</i> (D.Don) Soó	Narmada	Root, rhizome	Powder	Gojal	0.15	0.090909	1	11	MAS-094, 17
148	Orobanchaceae	<i>Pedicularis chelanthifolia</i> Schenk	Serfo spanthing	Leaf	Decoction	Skardu	0.3	0.166667	2	12	MAS-396, 12
149		<i>Pedicularis pectinatiformis</i> Bonati	Sunpo spanthing	Leaf	Infusion	Skardu	0.233333	0.157895	3	19	MAS-397, 12
150	Papaveraceae	<i>Corydalis crassifolia</i> Royle	Sackros/Zarvosh	Whole plant		Gojal	0.2	0.142857	1	7	MAS-093, 17
151		<i>Papaver somniferum</i> L.	Mardakhow	Latex	Decoction	Hunza	0.175	0.115385	3	26	MAS-136, 9, 33
152	Pinaceae	<i>Phus roxburghii</i> Sarg.	Chirpine	Resin, wood	Paste, powder, direct	Ghizer	0.233333	0.121212	4	33	MAS-047, 37
153		<i>Picea smithiana</i> (Wall.) Boiss.	Kachul	Resin, wood	Powder, decoction	Jalalabad	0.2	0.071429	1	14	MAS-201, 15
154		<i>Phus gerardiana</i> Wall. ex Lamb.	Cheenh	Resin, wood, leaf	Decoction, direct, paste, powder (K)	Jalalabad	0.175	0.096774	3	31	MAS-202, 15
155		<i>Phus wallichiana</i> A.B.Jacks.	Cheenh	Resin, wood, leaf	Infusion, powder	Nagar	0.175	0.142857	3	21	MAS-240, 15
156	Plantaginaceae	<i>Plantago ovata</i> Forsk.	Ispaghul	Seed, leaf, root	Powder, infusion	Ghizer	0.233333	0.069767	3	43	MAS-048, 16

Table 2 RFC, UV, number of uses, and ailments of species from each location (Continued)

S no.	Family	Species	Local name	Part used	Mode of use	Location	Average of RFC	Average of UV	Ailment categories	No. of use responses	Voucher no.	Previous citation
157		<i>Picrohiza kurroa</i> Royle ex Benth.	Karroo	Leaf, bark, root, rhizome	Paste	Kargha	0.2	0.130435	3	23	MAS-244	11
158		<i>Plantago major</i> L.	Shiltive, Boqna	Root, seed, leaf (K), seed (S), leaf, seed (Gh, H)	Direct, decoction, oil (S)	Ghizer, Hunza, Kargha, Skardu	0.2875	0.104482	9	127	MAS-073, MAS-161, MAS-282,	9, 14, 33, 37, 38, 53
159		<i>Plantago lanceolata</i> L.	Sman Hrswa, Sepilk, Yeeps	Flower, leaf (S), leaf, seed (H)	Decoction, infusion, paste (S), ash (H)	Gojal, Skardu	0.295833	0.121667	6	49	MAS-131, MAS-485	12, 17
160	Poaceae	<i>Zea mays</i> L.	Makayee	Fruit	Direct	Hunza	0.125	0.142857	1	7	MAS-139	9, 33
161		<i>Cymbopogon jwarancusa</i> (Jones) Schult.	Izkhari Makki	Flower	Decoction	Kargha	0.175	0.133333	2	15	MAS-242	40, 41
162		<i>Pennisetum glaucum</i> (L.) R.Br.	Chasq	Stem	Direct	Kargha	0.175	0.0625	1	16	MAS-243	39
163		<i>Saccharum bengalense</i> Retz.	Phoro	Root, stem	Powder	Kargha	0.175	0.190476	4	21	MAS-247	14
164		<i>Hordeum vulgare</i> L.	Cha Fay, York	Seed	Powder	Gojal, Kargha, Skardu	0.263889	0.161905	4	24	MAS-129, MAS-299,	17, 39
165		<i>Avena sativa</i> L.	Nas Choo, Sheshar	Seed (S), seed, leaf (H)	Decoction	Hunza, Skardu	0.270833	0.144796	5	47	MAS-190, MAS-496	9, 39
166	Polygonaceae	<i>Rheum tibeticum</i> Maxim. ex Hook. f.	Sheepod	Stem	Direct	Gojal	0.2	0.142857	1	7	MAS-102	17
167		<i>Bistorta amplexicaulis</i> (D.Don) Greene	Onbu	Root	Powder, decoction, infusion	Skardu	0.266667	0.214286	3	14	MAS-387	12
168		<i>Polygonum affine</i> D. Don.	Strin mindoq	Root, flower	Decoction, infusion	Skardu	0.4	0.142857	3	21	MAS-398	12
169		<i>Polygonum tataricum</i> L.	Bro Kho-Bro	Leaf, seed	Powder, decoction	Skardu	0.233333	0.157895	3	19	MAS-399	38
170		<i>Rheum spiciforme</i> Royle	Khakhol	Leaf, root	Direct, powder	Skardu	0.266667	0.214286	3	14	MAS-405	39
171		<i>Rumex chalepensis</i> Mill.	Sa-shing	Root	Decoction	Skardu	0.266667	0.083333	1	12	MAS-408	38
172		<i>Fagopyrum esculentum</i> Moench	Bro, Ghiawas, Stabro, Baraw	Seed (K, H, A), leaf, seed (S)	Direct, paste, powder	Astore, Hunza, Kargha, Skardu	0.195833	0.150985	12	89	MAS-016, MAS-146, MAS-259,	9, 13, 33, 38
173		<i>Rumex nepalensis</i> Spreng.	Chukkeen, Rashona	Root (K), leaf (A)	Paste	Astore, Kargha	0.229167	0.142857	2	14	MAS-021, MAS-263,	13, 14
174		<i>Rheum austrole</i> D.	Shoot, Lachu	Root (K, A), leaf, root	Powder, infusion,	Astore, Kargha, Skardu	0.280556	0.155556	8	50	MAS-030,	12, 13

Table 2 RFC, UV, number of uses, and ailments of species from each location (Continued)

S no.	Family	Species	Local name	Part used (Kargha-K, Nagar-N, Skardu-S, Ghizer-Gh, Hunza-H, Astore-A, Gilgit-G)	Mode of use (Kargha-K, Nagar-N, Skardu-S, Ghizer-Gh, Hunza-H, Astore-A, Gilgit-G)	Location (Astore-Astore, Ghizer-Ghizer, Gojal-Gojal, Central Hunza-Hunza, Jalalabad-Jalalabad, Kargha-Kargha, Nagar-Nagar, Skardu-Skardu)	Average of RFC	Average of UV	Ailment categories	No. of use responses	Voucher no.	Previous citation
175		<i>Oxyria digyna</i> (L.) Hill	Span Harswa, Skyurbutaq	Leaf (S), aerial (A)	Powder, decoction	Astore, Skardu	0.233333	0.177778	4	24	MAS-036, MAS-448	12, 38
176		<i>Rheum emodi</i>	Jarochnatal, Chontal	Whole plant	Decoction	Ghizer, Hunza	0.258333	0.120909	5	61	MAS-068, MAS-461	9, 16
177		<i>Rumex hastatus</i> D. Don	Churki	Whole plant (K), leaf, root, stem, fruit (Gn)	Direct, decoction, powder (Gn)	Ghizer, Kargha	0.3	0.093168	3	53	MAS-080, MAS-286	14, 16, 37
178		<i>Bistorta affinis</i> (D.Don) Greene	Buma	Leaf	Powder	Kargha, Skardu	0.204167	0.142857	2	14	MAS-507	39
179		<i>Polygonum hydropiper</i> L.	Thangmarcy	Leaf (K), aerial (S)	Decoction	Kargha, Skardu	0.2625	0.188235	5	27	MAS-333, MAS-518	14, 38
180	Primulaceae	<i>Primula macrophylla</i> D. Don	Benufsha	Whole plant	Decoction, powder	Gojal	0.175	0.083333	1	12	MAS-101	17
181		<i>Primula denticulata</i> Sm.	Daoo	Leaf, root	Decoction, powder, infusion (S)	Skardu	0.4	0.2	4	20	MAS-401	12
182		<i>Primula farinosa</i> L.	Spangpunar	Flower	Paste, decoction	Skardu	0.366667	0.142857	2	14	MAS-402	38
183	Ranunculaceae	<i>Clematis baicalensis</i> Qureshi & Chaudhri	Margush, Murgushi, Chinrik	Leaf, flower (H), whole plant (K)	Paste	Gojal, Jalalabad, Kargha	0.208333	0.086652	6	76	MAS-127, MAS-228, MAS-298	9, 14
184		<i>Aconitum repellicus</i> L.	Booma, Sai booma	Flower, leaf (N, H, K), whole plant (K), aerial (S)	Direct	Kargha, Nagar, Skardu	0.2	0.231481	7	31	MAS-316, MAS-378, MAS-501	11, 38, 53
185		<i>Ranunculus trichophyllum</i> Chaix ex Vill.	Threadleaf crowfoot	Whole plant	Paste, infusion	Kargha	0.175	0.111111	2	18	MAS-246	14
186		<i>Thalictrum foetidum</i> L.	Moneran	Leaf	Direct	Kargha	0.225	0.142857	1	7	MAS-248	39
187		<i>Aconitum violaceum</i> Jacquem. Ex. Stapf	Booma	Root	Decoction, powder	Skardu	0.233333	0.25	4	16	MAS-379	12
188		<i>Aquilegia fragrans</i> Benth.	Karfo Koo-kuk	Leaf, flower	Paste, decoction	Skardu	0.233333	0.133333	2	15	MAS-382	38
189		<i>Aquilegia pubiflora</i> Wall. Ex Royle	Koo-kuk	Leaf, flower	Paste	Skardu	0.233333	0.111111	2	18	MAS-383	38, 53
190		<i>Thalictrum foliolosum</i> DC.	Monyrun	Root	Decoction	Skardu	0.4	0.142857	2	14	MAS-412	38
191		<i>Delphinium brunonianum</i>	Makhoteng	Leaf, flower (K), whole plant (S, A)	Decoction, powder (S), infusion (S)	Astore, Kargha, Skardu	0.269444	0.132937	8	75	MAS-026, MAS-268, MAS-38, 39	12, 13,

Table 2 RFC, UV, number of uses, and ailments of species from each location (Continued)

S no.	Family	Species	Local name	Part used	Mode of use	Location (Astore-Astore, Ghizer-Ghizer, Gojal-Gojal, Central Hunza-Hunza, Jalalabad-Jalalabad, Kargha-Kargha, Nagar-Nagar, Skardu-Skardu)	Average of RFC	Average of UV	Ailment categories	No. of use responses	Voucher no.	Previous citation
192	Rosaceae	<i>Pulsatilla wallichiana</i> (Royle) Ulbr.	Zgiongmonana Looparimandoq	Flower	Powder	Astore, Skardu	0.316667	0.142857	2	14	MAS-038, MAS-450	
193	Rosaceae	<i>Comarum salesovianum</i> (Stephan) Asch. & Graebn.	Noghdurdom woodsh	Flower	Gojal		0.175	0.142857	1	7	MAS-092	9
194		<i>Potentilla eriocarpa</i> Wall. ex Lehmann.	Amber	Leaf, flower	Gojal		0.1	0.142857	1	7	MAS-099	17
195		<i>Potentilla microphylla</i> D. Don	Zatspinq	Leaf, seed	Gojal		0.175	0.222222	2	9	MAS-100	17
196		<i>Rubus irritans</i> Focke	Icheejeh	Fruit	Direct	Jalalabad	0.325	0.061224	3	49	MAS-207	15
197		<i>Potentilla argyrophylla</i> Wall. ex Lehmann.	Serfo Harswa	Whole plant	Paste	Skardu	0.466667	0.142857	1	7	MAS-400	12
198		<i>Rosa webbiana</i> Wall. ex Royle	Shighaye, Sia marpho, Chereer, Sia sarfo	Bark, wood (K), flower, bark (S), fruit, seed, wood (H, A)	Decoction	Astore, Gojal, Hunza, Jalalabad, Skardu	0.238333	0.103989	10	116	MAS-010, MAS-110, MAS-144, MAS-422	9, 13, 15, 17, 38, 53
199		<i>Prunus armeniaca</i> L.	Jui, Jaroty, Chooli	Fruit, kernel, oil	Direct, oil, powder, paste	Astore, Hunza, Jalalabad, Kargha, Nagar, Skardu	0.276389	0.082586	49	706	MAS-015, MAS-145, MAS-218, MAS-427	9, 13, 15, 32, 33, 38, 39
200		<i>Spiraea canescens</i> D.Don	Darah, Skhsri	Flower, stem, wood	Oil, decoction	Astore, Jalalabad	0.175	0.066667	3	52	MAS-018, MAS-219	13, 15
201		<i>Potentilla salessoviiana</i> Steph.	Sniarmastwa, Karfo mindoq	Flower	Infusion (S), paste	Astore, Kargha, Skardu	0.222222	0.116883	8	78	MAS-029, MAS-271, MAS-441	12, 13, 38
202		<i>Rosa brunonii</i> Lindl.	SiaMarpho, Sya	Bark	Decoction, powder (K), infusion (S)	Astore, Kargha, Skardu	0.216667	0.142857	3	21	MAS-031, MAS-273, MAS-443	12, 13
203		<i>Prunus dulcis</i> (Mill.) D.A.Webb	Badum, Balth, Kono, Stargi mar	Kernel, flower	Direct, oil, paste, decoction (S)	Ghizer, Hunza, Jalalabad, Kargha, Nagar, Skardu	0.231944	0.11893	21	194	MAS-069, MAS-223, MAS-334, MAS-462	15, 16, 32, 37, 38
204		<i>Rosa indica</i> L.	Ghulab	Flower	Paste, oil	Ghizer, Hunza, Skardu	0.313889	0.090149	6	62	MAS-076, MAS-164, MAS-469	
205		<i>Malus domestica</i> Borkh.	Skamkoosthu	Fruit	Powder	Kargha, Skardu	0.270833	0.142857	2	14	MAS-332, MAS-517	39
206		<i>Potentilla bifurca</i> L.	Tarqan	Flower (K), aerial (S)	Infusion (S), decoction (K)	Kargha, Skardu	0.345833	0.103343	4	54	MAS-334, MAS-519	12, 38, 39

Table 2 RFC, UV, number of uses, and ailments of species from each location (Continued)

S no.	Family	Species	Local name	Part used	Mode of use	Location (Astore-Astore, Ghizer-Ghizer, Gojal-Gojal, Central Hunza-Hunza, Jalalabad-Jalalabad, Kargha-Kargha, Nagar-Nagar, Skardu-Skardu)	Average of RFC	Average of UV	Ailment categories	No. of use responses	Voucher no.	Previous citation
207		<i>Prunus persica</i> (L.) Batsch	Takshu Choo	Fruit	Paste (K), decoction (S)	Kargha, Skardu	0.2375	0.171429	3	17	MAS-335, MAS-520	39, 53
208	Salicaceae	<i>Salix babylonica</i> L.	Muchhoor	Leaf, bark, seed, gum	Decoction, paste, direct	Hunza	0.125	0.142857	4	28	MAS-138	9
209		<i>Salix acmophylla</i> Boiss.	Brawoon	Leaf, bark, stem, branches	Decoction, paste	Jalalabad	0.275	0.074627	5	67	MAS-208	15, 53
210		<i>Salix denticulata</i> Andersson	Brawoon	Leaf, bark, stem, branches	Decoction, paste, direct	Jalalabad	0.225	0.083333	5	60	MAS-209	15
211		<i>Salix ilicifolia</i> Regel	Brawoon	Leaf, bark, stem, branches	Decoction, paste, direct	Jalalabad	0.175	0.09434	5	53	MAS-210	15
212		<i>Salix sericeocarpa</i> Andersson	Brawoon	Leaf, bark, stem, branches	Decoction, paste, direct	Jalalabad	0.175	0.096154	5	52	MAS-211	15
213		<i>Salix turanica</i> Nasarov	Brawoon	Leaf, bark, stem, branches	Decoction, paste, direct	Jalalabad	0.2	0.1	5	50	MAS-212	15
214		<i>Salix alba</i> L.	Mori Bayao, Bayo	Leaf, bark	Decoction, paste, direct	Ghizer, Kargha	0.225	0.099462	5	55	MAS-081, MAS-287	14, 16
215		<i>Populus alba</i> L.	Fulsoo, Turaq	Leaf, wood (K), leaf (H)	Decoction	Hunza, Jalalabad	0.1375	0.107143	2	21	MAS-172, MAS-231	9, 15, 33
216		<i>Populus nigra</i> L.	Jerpa	Leaf	Decoction	Hunza, Jalalabad, Nagar	0.166667	0.119048	3	28	MAS-175, MAS-234, MAS-363	15, 33
217		<i>Salix tetrasperma</i> Roxb.	Byao, Bew	Leaf, bark	Decoction, paste, direct	Hunza, Kargha	0.2	0.107143	2	21	MAS-177, MAS-303	9, 14
218	Saxifragaceae	<i>Saxifraga hirculus</i> L.	Sitbark	Whole plant	Decoction	Gojal	0.15	0.117647	4	34	MAS-103	17
219		<i>Bergenia himalaica</i> Boniss.	Sanspur	Root	Powder, decoction	Kargha	0.175	0.133333	2	15	MAS-241	9, 11, 12, 16, 33, 53
220		<i>Bergenia stracheyi</i> (Hook.f. & Thomson) Engl.	Sasper, Khichlay	Root, leaf	Infusion, powder, paste, direct	Ghizer, Hunza, Nagar, Skardu	0.25625	0.14944	18	124	MAS-075, MAS-163, MAS-357, MAS-468	9, 11, 12, 16, 33
221		<i>Bergenia ciliata</i> (Haw.) Steimb.	Shafus, Shaphus	Leaf (K), leaf, seed (S)	Powder, decoction, direct	Kargha, Skardu	0.308333	0.162338	12	75	MAS-321, MAS-506	12, 13, 39
222	Solanaceae	<i>Solanum nigrum</i> L.	Gabeeli, Gabilo, Drumbashokhlo	Whole plant (N/Gh), leaf, fruit (K), fruit, seed (H), fruit (A)	Direct, decoction (K), powder (Gh)	Astore, Ghizer, Gojal, Kargha, Nagar	0.243333	0.125426	20	190	MAS-005, MAS-053, MAS-252, MAS-343	13, 14, 16, 17, 37
223		<i>Hyoscyamus niger</i> L.	Landlungstwa	Seed	Paste, poultice, decoction	Astore, Kargha, Skardu	0.25	0.15873	3	19	MAS-027, MAS-269, MAS-439	13, 53
224		<i>Datura stramonium</i> L.	Datuvo, Datura	Seed (H), flower, fruit, seed, leaf (Gh, K)	Decoction, ash (K), paste (K)	Ghizer, Hunza, Kargha	0.266667	0.105286	13	138	MAS-071, MAS-159,	9, 14, 16, 33,

Table 2 RFC, UV, number of uses, and ailments of species from each location (Continued)

S no.	Family	Species	Local name	Part used (Kargha-K, Nagar-N, Skardu-S, Ghizer-Gh, Hunza-H, Astore-A, Gilgit-G)	Mode of use (Kargha-K, Nagar-N, Skardu-S, Ghizer-Gh, Hunza-H, Astore-A, Gilgit-G)	Location (Astore-Astore, Ghizer-Ghizer, Gojal-Gojal, Central Hunza-Hunza, Jalalabad-Jalalabad, Kargha-Kargha, Nagar-Nagar, Skardu-Skardu)	Average of RFC	Average of UV	Ailment categories	No. of use responses	Voucher no.	Previous citation
225		<i>Capsicum annuum</i> L.	Marooch	Fruit	Direct	Hunza, Nagar	0.2375	0.065613	5	101	MAS-187, 9 MAS-370	
226	Tamaricaceae	<i>Myricaria squamosa</i> Desv.	Targ	Leaf, flower	Powder	Gojal	0.175	0.178571	5	28	MAS-097 17	
227	Thymelaeaceae	<i>Daphne mucronata</i> Royle	Nirko	Leaf, fruit, wood	Paste, poultice	Jalalabad	0.275	0.088889	4	45	MAS-193 15	
228	Urticaceae	<i>Urtica dioica</i> L.	Khaeshing	Whole plant (S, K), leaf, root (Gh)	Direct, decoction, paste (S)	Ghizer, Kargha, Skardu	0.236111	0.132675	12	149	MAS-087, 13, 37, MAS-293, 53 MAS-475	
229	Violaceae	<i>Viola serpens</i> Wall. ex Ging.	Skora mindoq, Lillo	Flower (S), whole plant (Gh, H)	Decoction	Ghizer, Hunza, Skardu	0.258333	0.124008	6	58	MAS-077, 9, 16, MAS-165, 33, 38 MAS-470	
230	Zingiberaceae	<i>Curcuma longa</i> L.	Halichi	Stem	Powder	Hunza, Nagar	0.275	0.088933	4	45	MAS-188, 32 MAS-371	
231	Zygophyllaceae	<i>Tribulus terrestris</i> L.	Kokoloq, Kokoring, Hulk ga kurice	Whole plant (Gh, K, A), seed (S), fruit (H)	Paste, decoction	Astore, Ghizer, Hunza, Kargha, Skardu	0.23	0.142857	9	63	MAS-008, 9, 13, MAS-056, 33, 37, MAS-255, 38, 39 MAS-420	




Fig. 8 Percentage of the parts of plants used for medicinal purposes

led many local people to over-exploit this valuable resource. Over-exploitation of medicinal plant species is widespread across the region, exacerbated by some local people attempting to maximize financial benefits in a single harvest, with little concern for the ramifications for subsequent years [9, 30, 34, 42].

This study clearly reveals the importance and contributions of the THPs and retailers as well as the transfer of knowledge within the families from elders to the younger generation for the retention of indigenous medicinal knowledge, where and when to acquire a particular species and the utilization of medicinal plants. The THPs preserved existing knowledge and showed a great deal of openness to knowledge sharing. They attributed the loss of knowledge between generations, not to any failure of their own to impart knowledge, but rather to a lack of learning aptitude in the younger generation [34, 106]. Retailers have adapted well to the demands of different ethnic and tribal groups. These factors point towards a high level of cooperation, collaboration, and openness to knowledge exchange amongst the ethnicities and tribes of Gilgit-Baltistan. Local public and private institutes can therefore play a vital role in clustering the knowledge and bridging the gaps by providing platforms for recording, sharing, and disseminating traditional knowledge.

We found that Gilgit-Baltistan's position as a gateway between the Central and South Asia caused its exposure to a number of traditional medicinal systems including the Ayurveda, traditional Chinese medicine, Unani, and Tibetan—which highly influenced traditional medicine knowledge in this region [3, 46]. Our study design and timeline restricted us from further exploration of these historical details. Therefore, we were not able to explore the timeline and actual contributions of these systems to local knowledge. However, it appears likely that medicinal practices in Hunza, Nagar, Ghizer, and Gilgit were influenced by traditional Chinese medicine (TCM) system while the remaining part of the study area was dominated by a mix of Ayurveda and Unani systems. This is an interesting finding and deserves further research. Most parts of Pakistan are primarily relying on a mix of Unani and Ayurveda medicinal systems—a combination which is rarely found elsewhere [107]. A dedicated study exploring the approaches followed by these medicinal systems, their complementarities, and differences could lead to the generation of highly valuable scientific findings that could contribute to the communities relying on these systems globally.

With the involvement of multiple stakeholders (the relevant local government departments, herbal medicine-producing companies, THPs, and the interest of the national

government), medicinal plants and associated traditional knowledge from Gilgit-Baltistan can make a substantial contribution to traditional health practices at a national level as well as contribute significantly to the national market and the livelihood resources of local communities. Proper licensing will allow the THPs to legally practice, document, and disseminate their knowledge. The concerned government departments can provide a platform for THPs from the region to get registered and licensed as hakims [39]. Our effort to involve school students in the collection of data is a way of exposing the younger generation to identification of their resources and developing their interest in traditional knowledge and why it is important to ensure its transfer to them from the older generation. Such a consortium will also prove beneficial for the production of medicinal plants on a commercial scale, their sustainable utilization, and organizing refreshers on different aspects associated with medicinal plant resources for the local THPs, retailers, and collectors in order to ensure an optimal and efficient utilization of the available resources.

These points are of utmost importance when it comes to the conservation and transfer of traditional medicinal knowledge to future generations. Worldwide, patients are increasingly opting for medications involving traditional techniques, herbal medicine, and meditation [108, 109]. Gilgit-Baltistan has natural medicinal resources, a vast indigenous knowledge bank, and most importantly one of the best mountainous landscapes for tourism and meditation. It is doubtless the best option to be considered for developing into a sanctuary through government interventions. Gilgit-Baltistan, considered to be home to the ideology of "SHANGRI LA" with its abundant natural resources, can provide a home to those who seek medication through centuries-old traditional knowledge, sacredly transferred from one generation to another.

Conclusions and recommendations

The diverse plant resources and the geographical importance of the region for trade and travel routes, historically made Gilgit-Baltistan a hotspot for cultural, religious, and traditional knowledge exchange. Being part of an ancient trade route, the resident communities adapted and upgraded their traditional healing systems through interactions with the Indian subcontinent, China, Scythia, Transoxiana, and Ancient Greece. This influence and amalgamation of Chinese, Ayurveda, Unani, and Tibetan medicinal systems is apparent in local traditional knowledge. Our study revealed that most of the local people still rely on indigenous healing practices. Higher knowledge and use of medicinal plants is retained in the areas that also serve as main trade centers in the region. The trade of medicinal plants in the region is the one key factor in

retaining traditional knowledge on medicinal plant utilization. This continued reliance on medicinal plants shows the significance of these traditional practices. A thorough evaluation is needed by ethno-pharmacologists and other concerned institutions working for public health and hygiene, especially focusing on THPs, market actors and old folk from the region. For strategies to be devised for market exploration, raising awareness, and continuity of TK, involvement is required from Government institutions, research organizations, NGOs, donors, and the private sector.

Additional files

Additional file 1: Primary data on medicinal plants and their uses collected during the field survey. (XLSX 636 kb)

Additional file 2: RFC and UV of species reported from Central Hunza. (XLSX 14 kb)

Additional file 3: RFC and UV of species reported from Ghizer. (XLSX 14 kb)

Additional file 4: RFC and UV of species reported from Gojal Hunza. (XLSX 13 kb)

Additional file 5: RFC and UV of species reported from Jalalabad. (XLSX 13 kb)

Additional file 6: RFC and UV of species reported from Kargha. (XLSX 17 kb)

Additional file 7: RFC and UV of species reported from Nagar. (XLSX 13 kb)

Additional file 8: RFC and UV of species reported from Skardu. (XLSX 19 kb)

Additional file 9: RFC and UV of species reported from Astore. (XLSX 13 kb)

Additional file 10: Top 10 species from each location for number of uses, ailment categories, RFC, and UV. (XLSX 14 kb)

Additional file 11: List of publications from the region reporting on medicinal plants and their uses. (XLSX 12 kb)

Additional file 12: Ailment categories. (XLSX 14 kb)

Abbreviations

CPEC: China-Pakistan Economic Corridor; DA: Discriminant analysis; FC: Frequency of citation; FGD: Focused group discussion; HH: Household; HKH: Himalaya Karakoram Hindukush Mountain Range; ICF: Informant consensus factor; KKH: Karakoram Highway; MAPs: Medicinal and aromatic plants; NGO: Non-Governmental Organization; RFC: Relative frequency of citation; TCM: Traditional Chinese medicine; THP: Traditional health practitioner; TK: Traditional knowledge; UV: Use value

Acknowledgements

We are thankful to the participants of the field survey for their cooperation and time. The Law Enforcement Agencies at Gilgit-Baltistan especially Captain Faisal Anwar, Captain Muhammad Awais Salim, and their entire team are thanked for their support and provision of services in times of need.

Funding

The study was conducted under the "Investigating and recalibrating traditional ecological calendars and climate change adaptation in the Himalaya region" funded by Bureau of International Cooperation Chinese Academy of Sciences, Grant No. 151853KYSB20160032. Additional support was provided by the National Natural Science Foundation of China, Grant No. 41661144001. Funds acquired from these sources were utilized for field visits, collecting, and analyzing data and interpretation of results.

Availability of data and materials

In addition to the data presented through tables and figures in the main text, all the data supporting the findings and results is available in the additional files provided with the manuscript.

Authors' contributions

MAS, SR, and JX conceptualized, planned, and designed the study. MAS and TK led the data collection team of SA, CK, AP, ZB, and SB. MAS and SR analyzed the data, wrote, and finalized the manuscript. RH and JX revised the paper. SR and JX supervised the first author during his doctoral research. All the authors have read and approved the final manuscript prior to submission.

Ethics approval and consent to participate

The study is based on field surveys, and no human or animal trials were involved or conducted. Formal consent was obtained from participants prior to data collection and publication. In addition, International Society of Ethnobiology (ISE) code of ethics <http://www.ethnobiology.net> were strictly followed while conducting the research.

Consent for publication

Formal consent was obtained from participants prior to data collection and publication.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Key Laboratory for Plant Diversity and Biogeography of East Asia, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, Yunnan, China. ²University of Chinese Academy of Sciences, Beijing 100049, China. ³World Agroforestry Centre (ICRAF), East and Central Asia Office, Kunming 650201, Yunnan, China. ⁴Missouri Botanical Garden, Post Office Box 299, St. Louis, MO 63166, USA. ⁵Department of Biological Sciences, Karakorum International University, Gilgit, Gilgit-Baltistan, Pakistan.

Received: 31 August 2018 Accepted: 9 December 2018

Published online: 28 January 2019

References

- Biddulph J. Tribes of the Hindoo Koosh. Graz: Akademische Druck- u. Verlagsanstalt; 1971. Available from: file: <https://catalog.hathitrust.org/Record/100159287>.
- Khan T, Khan IA, Rehman A, Karim R, Bibi S, Rukhsana. Investigation into ethnopharmacological perceptions and practices among traditional communities of Karakorum Mountain Ranges: special reference to Berberis species. Asian J Biol Life Sci. 2014;3:173–8.
- Hauptmann H. Pre-islamic heritage in the northern areas of Pakistan. In: Bianca S, editor. Karakoram Hidden treasures North Areas Pakistan. 2nd ed. Turin: Umberto Allemandi & Co.; 2007. p. 21–40.
- Irshad MS, Xin Q, Arshad H. One belt and one road: dose China-Pakistan Economic Corridor benefit for Pakistan's economy? J Econ Sustain Dev. 2015;6:200–7 Available from: <http://ssrn.com/abstract=2710352>.
- Ullah S, Muhammad A, Hussain I, Hafeez-Ur-Rehman, Hyder MZ. Geneticalysis of economically important apricot cultivars in Gilgit Baltistan based on SSR molecular markers. Rom Biotechnol Lett. 2017;22:12456–63.
- Ishaq S, Khan MZ, Begum F, Hussain K, Amir R, Hussain A, et al. Climate change impact on mountain biodiversity: a special reference to Gilgit-Baltistan of Pakistan. J Mt Area Res. 2015;1:53–63.
- Khan T, Rukhsana. Online earning, a promising source impacting poverty and women empowerment: a case study from high mountainous nexus of Himalaya-Karakoram-Hindukush, Pakistan. Nat Sci. 2016;14:12–6.
- Joshi S, Jasra WA, Ismail M, Shrestha RM, Yi SL, Wu N. Herders' perceptions of and responses to climate change in Northern Pakistan. Environ Manag. 2013;52:639–48.
- Bano A, Ahmad M, ben Hadda T, Saboor A, Sultana S, Zafar M, et al. Quantitative ethnomedicinal study of plants used in the Skardu valley at high altitude of Karakoram-Himalayan range, Pakistan. J Ethnobiol Ethnomed. 2014;10:43 Available from: <http://ethnobiomed.biomedcentral.com/articles/10.1186/1746-4269-10-43>.
- Abbas A, Khan M, Shah MH, Shah M, Pervez A, Ahmad M. Ethnobotanical appraisal and cultural values of medicinally important wild edible vegetables of Lesser Himalayas-Pakistan. J Ethnobiol Ethnomed. 2013;9:66 Available from: <http://ethnobiomed.biomedcentral.com/articles/10.1186/1746-4269-9-66>.
- Khan T, Khan IA, Rehman A, Ali H. Ethnobotanical studies on non-medicinal plants of Shinaki Valley Hunza, Gilgit-Baltistan. Int J Biosci. 2013;3:63–70 Available from: http://www.researchgate.net/profile/Haibat_Ali/publication/264534933_Ethnobotanical_studies_on_non-medicinal_plants_of_Shinaki_Valley_Hunza_Gilgit-Baltistan/links/53e33d190cf2b9d0d8331867.pdf.
- Khan SW, Khatoon S. Ethnobotanical studies on useful trees and shrubs of Haramosh and Bugrote valleys in Gilgit northern areas of Pakistan. Pakistan J Bot. 2007;39:699–710 Available from: [http://www.pakbs.org/pjbot/PDFs/39\(3\)/PJB39\(3\)699.pdf](http://www.pakbs.org/pjbot/PDFs/39(3)/PJB39(3)699.pdf).
- Shah F, Asghari B. Ethnobotanical and physiological studies of some endangered plant species collected from two different altitudes in Gilgit Baltistan. Pakistan J Bot. 2012;44:165–70.
- Khan B, Abdukadir A, Qureshi R, Mustafa G. Medicinal uses of plants by the inhabitants of Khunjerab national park, Gilgit, Pakistan. Pakistan J Bot. 2011; 43:2301–10.
- Khan SW, Khatoon S, Abbas Q, Raza G, Hussain A. Inventory of the alpine flora of Haramosh and Bagrote valleys (Karakoram range) district Gilgit, Gilgit-Baltistan, Pakistan. Pakistan J Bot. 2016;48:1559–72.
- Hussain I, Bano A, Ullah F. Traditional drug therapies from various medicinal plants of Central Karakoram National Park, Gilgit-Baltistan, Pakistan. Pakistan J Bot. 2011;43:79–84.
- Khan T, Khan IA, Rehman A. Assessment of ethno-ecological knowledge, practices and perceptions among traditional communities of Karakorum Mountain Ranges: a special reference to Berberis spp. J Biodivers Environ Sci. 2014;5:258–66.
- Khan SW, Abbas Q, ul Hassan SN, Khan H, Hussain A. Medicinal Plants of Turmic Valley (Central Karakoram National Park), Gilgit-Baltistan, Pakistan. J Bioresour Manag. 2015;2 Available from: <http://corescholar.libraries.wright.edu/jbm>.
- Khan T, Khan IA, Rehman A. A composite review of Berberis in service of humanity-in-trouble: an ethnopharmacological and pharmacognosic perspective. J Biodivers Environ Sci. 2015;6:21–33.
- Sheng-Ji P. Ethnobotanical approaches of traditional medicine studies: some experiences from Asia. Pharm Biol. 2001;39:74–9 Available from: <http://www.tandfonline.com/doi/full/10.1076/phbi.39.s1.74.0005>.
- Mugisha MK, Asiimwe S, Namutebi A, Borg-Karlson AK, Kakudidi EK. Ethnobotanical study of indigenous knowledge on medicinal and nutritious plants used to manage opportunistic infections associated with HIV/AIDS in western Uganda. J Ethnopharmacol. 2014;155:194–202. <https://doi.org/10.1016/j.jep.2014.05.012> Elsevier.
- Zhang L, Zhuang H, Zhang Y, Wang L, Zhang Y, Geng Y, et al. Plants for health: an ethnobotanical 25-year repeat survey of traditional medicine sold in a major marketplace in North-west Yunnan, China. J Ethnopharmacol. 2018;224: 119–25. <https://doi.org/10.1016/j.jep.2018.05.029> Elsevier Ireland Ltd.
- Nesheim I, Dhillon SS, Stølen KA. What happens to traditional knowledge and use of natural resources when people migrate? Hum Ecol. 2006;34:99–131.
- Ju Y, Zhuo J, Liu B, Long C. Eating from the wild: diversity of wild edible plants used by Tibetans in Shangri-la region, Yunnan, China. J Ethnobiol Ethnomed. 2013;9:28.
- Indigenous peoples and climate change. In: Salick J, Byg A, editors. Indig. Peoples Clim. Chang. Oxford: Tyndall Centre for Climate Change Research, 2007. Available from: <http://ir.lib.uwo.ca/iipj/vol1/iss1/2>.
- Zeven AC, Wet J m. J De. Dictionary of cultivated plants and their centres of diversity: Excluding most ornamentals, forest trees and lower plants. 2nd ed. Dict. Cultiv. plants their centres Divers. Wageningen: Centre for Agricultural Publishing and Documentation, Wageningen; 1982.
- Khan T, Ali S, Khan IA, Alam J. Growth responses of *Hippophae rhamnoides* L. subsp. sinensis towards different agro-climatic conditions. Int J Biosci. 2013;3:293–7 Available from: <http://www.innspub.net/wp-content/uploads/2013/09/JB-V3No9-p293-297.pdf>.
- Yaseen G, Ahmad M, Sultana S, Suleiman Alharrasi A, Hussain J, Zafar M, et al. Ethnobotany of medicinal plants in the Thar Desert (Sindh) of Pakistan. J Ethnopharmacol. 2015;163:43–59.
- Khan Shinwari Z, Watanabe T, Ali M, Anwar R. Medicinal plants in 21st century. In: International symposium on medicinal linkages beyond national boundaries. Kohat: Kohat University of Science and Technology; 2005. p. 12–7.
- Shedai A, Gulshan B. Ethnomedicinal uses of plant resources in Gilgit-Baltistan of Pakistan. J Med Plants Res. 2012;6:4540–9.
- Shinwari ZK, Qaiser M. Efforts on conservation and sustainable use of medicinal plants of Pakistan. Pakistan J Bot. 2011;43:5–10.

32. Rizvi MA. Elemental composition of medicinal flowers. *Pakistan J Bot.* 2007; 39:2541–52.
33. Khan T, Khan IA, Rehman A, Alam J, Ali S. Exploration of near-extinct folk wisdom on medicinally important plants from Shinaki Valley Hunza, Pakistan. *Int J Biosci.* 2013;3:180–6. Available from: <https://www.cabdirect.org/cabdirect/abstract/20133395619>.
34. Shedad AA, Xu M, Gulraiz B. Traditional medicinal uses of plants in Gilgit-Baltistan, Pakistan. *J Med Plant Res.* 2014;8:992–1004 Available from: <http://academicjournals.org/journal/JMPR/article-abstract/B2A4EA046850>.
35. Abbas Z, Khan SM, Abbasi AM, Pieroni A, Ullah Z, Iqbal M, et al. Ethnobotany of the Balti community, Tormik valley, Karakorum range, Baltistan, Pakistan. *J Ethnobiol Ethnomed.* 2016;12:38 Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27612599%5Cn> <http://www.ncbi.nlm.nih.gov/articlerender.fcgi?artid=PMC5018187>.
36. Akhter S, Akhtar N, Kazin S, Khan T. Study of ethno-gynecologically important medicinal and other plants used for women specific purposes in Murazaabadi, Hunza, Pakistan. *Nat Sci.* 2016;14:36–9.
37. Ismail M, Haq N, Ali S, Ali I, Shaheen R, Murtaza G, et al. Qualitative analysis for phytochemicals of selected medicinal plants from Gilgit-Baltistan, Pakistan. *Asian J Chem.* 2017;29:1929–32 Available from: http://www.asianjournalofchemistry.co.in/User/ViewFreeArticle.aspx?ArticleID=29_9_12.
38. Ali I, Hussain H, Batool H, Dad A, Raza G, Falodun A, et al. Documentation of ethno veterinary practices in the CKNP region, Gilgit-Baltistan. *Int J Phytomedicine.* 2017;9:223–40.
39. Shinwari ZK, Gilani SS. Sustainable harvest of medicinal plants at Bulashbar Nullah, Astore. *J Ethnopharmacol.* 2003;84:289–98.
40. Athar M, Siddiqi MA. Reflections on the taxonomy and distribution of medicinal flowers of Pakistan. *SIDA Contrib to Bot.* 2004;21:357–68.
41. Qureshi RA, Ghufran MA, Sultana KN, Ashraf M, Khan AG. Ethnobotanical studies of medicinal plants of Gilgit District and surrounding areas. *Ethnobot Res Appl.* 2006;5:115–22.
42. Kayani S, Ahmad M, Sultana S, Khan Shinwari Z, Zafar M, Yaseen G, et al. Ethnobotany of medicinal plants among the communities of alpine and sub-alpine regions of Pakistan. *J Ethnopharmacol.* 2015;164:186–202. <https://doi.org/10.1016/j.jep.2015.02.004> Elsevier.
43. Khan T, Khan IA, Rehman A, Bibi Z. Determination of effectiveness of berberine, a characteristic phytochemical of Berberis species, against human proteome using in-silico analysis. *J Biodivers Environ Sci.* 2014;4:53–63 Available from: <http://innspub.net/wp-content/uploads/2014/05/JBES-Vol4No5-p53-63.pdf>.
44. Khan T, Khan IA, Rehman A. Assessment of geographic distribution and taxonomic anomalies of Berberis species found and reported from Karakoram, Hindukush and Western Himalayan Mountain Ranges in Pakistan. *J Biodivers Environ Sci.* 2014;5:267–73.
45. Abbas Z, KHAN SM, Alam J, Ullah Z, Khan SW, Alam N. Species diversity and phyto-climatic gradient of a montane ecosystem in the Karakorum range. *Pakistan J Bot.* 2017;49:89–98.
46. Barsha SBAK. Aks-e-Gilgit Baltistan. Gilgit: North Books Gilgit; 2013.
47. Stevens CJ, Murphy C, Roberts R, Lucas L, Silva F, Fuller DQ. Between China and South Asia: a middle Asian corridor of crop dispersal and agricultural innovation in the Bronze Age. *The Holocene.* 2016;26: 1541–55.
48. Middleton R. The great game – myth or reality? Tajikistan High Pamirs. 2011; Available from: http://pamirs.org/ExtractOdyssey_Great_Game.pdf.
49. Kassam K, Karamkhudeva M, Ruelle M, Baumflel M. Medicinal plant use and health sovereignty: findings from the Tajik and Afghan Pamirs. *Hum Ecol.* 2010;38:817–29.
50. Mark JJ. Silk road. 2014 [Cited 27 Mar 2018]. Available from: https://www.anciente.eu/Silk_Road/
51. Attewell G. Compromised: making institutions and indigenous medicine in Mysore State, circa 1908–1940. *Cult Med Psychiatry.* 2014;38:369–86.
52. Introduction. 2016 [Cited 30 Mar 2018]. Available from: <http://ayush.gov.in/about-the-systems/unani/introduction>
53. Wazir SM, Dasti AA, Shah J. Common medicinal plants of Chapursan Valley, Gojal II, Gilgit-Pakistan. *J Res (Science) Bahauddin Zakariya Univ Multan Pakistan.* 2004;15:41–3.
54. Khan T, Khan IA, Rehman A. A review on Berberis species reported from Gilgit-Baltistan and Central Karakoram National Park, Pakistan. *J Med Plants Stud.* 2014;2:16–20.
55. Muhammad G. Festivals and folklore of Gilgit. Culcutta: The Asiatic Society, 57, Park Street; 1905.
56. Geography and demography of Gilgit Baltistan. [cited 18 July 2017]. Available from: <http://www.gilgiltbastanscouts.gov.pk/geodemo.htm>
57. Khan T, Khan IA, Rehman A. Investigations into ecological consequences and threats from ethnoecological and ethnobotanical practices across Karakorum Mountain Ranges: a case study Berberis. *Curr World Environ.* 2014;9:713–20 Available from: <http://www.cwejournal.org/vol9no3/investigations-in-to-ecological-consequences-and-threats-from-ethnoecological-and-ethnobotanical-practices-across-karakorum-mountain-ranges-a-case-study-berberis>.
58. Rhoades RE, Thompson SI. Adaptive strategies in alpine environments: beyond ecological particularism. *Am Ethnol.* 1975;2:535–51.
59. Kreutzmann H. Challenge and response in the Karakoram: socioeconomic transformation in Hunza, Northern Areas, Pakistan. *Mt Res Dev.* 1993;13:19–39.
60. IFAD. Economic transformation initiative Gilgit-Baltistan. 2015. Available from: <https://webapps.ifad.org/members/eb/114/docs/EB-2015-114-R-14-Project-design-report.pdf>
61. Platon A. Pakistan ethnic map. 2012. Available from: https://commons.wikimedia.org/wiki/File:Pakistan_ethnic_map.svg. Cited 10 Apr 2018
62. Khan T, Khan IA, Rehman A, Karim R, Rukhsana, Ali S. Evaluation of human induced threats on berberis populations across cultural landscape of Karakorum Mountain Ranges. *J Biodivers Environ Sci.* 2014;5:333–42.
63. Khan T, Khan IA, Rehman A, Karim R, Rukhsana. Studies on population change dynamics of critically endanger Berberis species in Karakoram Mountain Ranges: an ethnoecological perspective. *Asian J Biol Life Sci.* 2014; 3:206–11.
64. Gautam AP, Webb EL, Eiumnoph A. GIS assessment of land use/land cover changes associated with community forestry implementation in the middle hills of Nepal. *Mt Res Dev.* 2002;22:63–9 International Mountain Society. Available from: [https://doi.org/10.1659/0276-4741\(2002\)022\[0063:GAOLUL\]2.0.CO;2](https://doi.org/10.1659/0276-4741(2002)022[0063:GAOLUL]2.0.CO;2).
65. Maikhuri RK, Rao KS, Saxena KG. Bioprospecting of wild edibles for rural development in the Central Himalayan Mountains of India. *Mt Res Dev.* 2004;24:110–3 Available from: <http://www.scopus.com/inward/record.url?eid=2-s2.0-3042813290&partnerID=40&md5=02e54e5d71a4ab7ac914054b2f87471>.
66. Liniger H, Schwilch G. Enhanced decision-making based on local knowledge. *Mt Res Dev.* 2002;22:14–8 Available from: <http://www.bioone.org/doi/pdf/10.1659/0276-4741%282002%29022%5B0014%6AEDMBOL%5D2.0.CO;2>.
67. Khan T, Khan IA, Rehman A, Karim R. Conserving wildlife, not their clinics: a case study of wild herbal clinic (Berberis) across alpine pastures in Karakoram Mountain Ranges. *Asian J Biol Life Sci.* 2014;3(3):195–9.
68. AKRSP, JICA. Basic study on horticulture sector in Gilgit-Baltistan. 2010.
69. Gratani M, Bohensky EL, Butler JRA, Sutton SG, Foale S. Experts' perspectives on the integration of indigenous knowledge and science in wet tropics natural resource management. *Aust Geogr.* 2014;45:167–84.
70. Gazzaneo LRS, Paiva de Lucena RF, de Albuquerque UP. Knowledge and use of medicinal plants by local specialists in an region of Atlantic Forest in the state of Pernambuco (Northeastern Brazil). *J Ethnobiol Ethnomed.* 2005;1:1–8.
71. Bradacs G, Heilmann J, Weckerle CS. Medicinal plant use in Vanuatu: a comparative ethnobotanical study of three islands. *J Ethnopharmacol.* 2011; 137:434–48. <https://doi.org/10.1016/j.jep.2011.05.050> Elsevier Ireland Ltd.
72. Ladio AH, Molares S. The dynamics of use of nontraditional ethnobiological products: some aspects of study. In: Albuquerque UP, da Cunha LVFC, de Lucena RFP, Alves RRN, editors. *Methods Tech Ethnobiol ethnoecology.* New York: Springer International Publishing; 2014. p. 311–9.
73. Sher H, Bussmann RW, Hart R, De Boer HJ. Traditional use of medicinal plants among Kalasha, Ismaeli and Sunni groups in Chitral District, Khyber Pakhtunkhwa province, Pakistan. *J Ethnopharmacol.* 2016;188:57–69. <https://doi.org/10.1016/j.jep.2016.04.059>. Elsevier.
74. Bernard HR. Sampling III: nonprobability samples and choosing informants. *Res Methods Anthropol Qual Quant Approaches.* 2011;143–55. 5th Edition. Altamira press.
75. Ranjiktar S, Rajbhandary S. Utilization of medicinal plants as home herbal-remedy in some urban areas of Kathmandu. *Nepal J Plant Sci.* 2008;2:56–65.
76. Benarba B. Medicinal plants used by traditional healers from South-west Algeria: an ethnobotanical study. *J Intercult Ethnopharmacol.* 2016;5:320 Available from: <http://www.scopemed.org/fulltextpdf.php?mno=228905>.

77. Teklehaymanot T. Ethnobotanical study of knowledge and medicinal plants use by the people in Dek Island in Ethiopia. *J Ethnopharmacol.* 2009;124:69–78. <https://doi.org/10.1016/j.jep.2009.04.005> Elsevier Ireland Ltd.
78. Umair M, Altaf M, Abbasi AM. An ethnobotanical survey of indigenous medicinal plants in Hafizabad district, Punjab-Pakistan. *PLoS One.* 2017;12:1–22. <https://doi.org/10.1371/journal.pone.0177912>.
79. Singh AG, Kumar A, Tewari DD. An ethnobotanical survey of medicinal plants used in Terai forest of western Nepal. *J Ethnobiol Ethnomed.* 2012;8:14 Available from: <https://ethnobiomed.biomedcentral.com/articles/10.1186/1746-4269-8-19>.
80. Shinwari S, Ahmad M, Luo Y, Zaman W. Quantitative analyses of medicinal plants consumption among the inhabitants of Shangla-Kohistan areas in Northern-Pakistan. *Pakistan J Bot.* 2017;49:725–34.
81. Georgakarakos S, Kitsiou D. Mapping abundance distribution of small pelagic species applying hydroacoustics and Co-Kriging techniques. *Hydrobiologia.* 2008;612:155–69.
82. Kanankege KST, Alkhamis MA, Phelps NBD, Perez AM. A probability Co-Kriging model to account for reporting bias and recognize areas at high risk for Zebra Mussels and Eurasian Watermilfoil invasions in Minnesota. *Front Vet Sci.* 2018;4:1–10 Available from: <http://journal.frontiersin.org/article/10.3389/fvets.2017.00231/full>.
83. Hernández-Stefanoni JL, Alberto Gallardo-Cruz J, Meave JA, Dupuy JM. Combining geostatistical models and remotely sensed data to improve tropical tree richness mapping. *Ecol Indic.* 2011;11:1046–56.
84. Trotter R, Logan M. Informant consensus: a new approach for identifying potentially effective medicinal plants. In: Etkin NL, editor. *Plants Indig Med Diet Behav Approaches;* 1986. p. 91–112.
85. Amjad MS, Qaeem MF, Ahmad I, Khan SU, Chaudhari SK, Malik NZ, et al. Descriptive study of plant resources in the context of the ethnomedicinal relevance of indigenous flora: a case study from Toli Peer National Park, Azad Jammu and Kashmir, Pakistan. *PLoS One.* 2017;12:1–31.
86. Heinrich M, Edwards S, Moerman DE, Leonti M. Ethnopharmacological field studies: a critical assessment of their conceptual basis and methods. *J Ethnopharmacol.* 2009;124:1–17. <https://doi.org/10.1016/j.jep.2009.03.043> Elsevier Ireland Ltd.
87. Rehman MN, Ahmad M, Sultana S, Zafar M, Edwards S. Relative popularity level of medicinal plants in Talagang, Punjab Province, Pakistan. *Brazilian J Pharmacogn.* 2017;27:751–75 Available from: <https://www.sciencedirect.com/science/article/pii/S0102695X17302053>.
88. Butt MA, Ahmad M, Fatima A, Sultana S, Zafar M, Yaseen G, et al. Ethnomedicinal uses of plants for the treatment of snake and scorpion bite in Northern Pakistan. *J Ethnopharmacol.* 2015;168:164–81. <https://doi.org/10.1016/j.jep.2015.03.045> Elsevier.
89. Vitalini S, Iriti M, Puricelli C, Ciuchi D, Segale A, Fico G. Traditional knowledge on medicinal and food plants used in Val San Giacomo (Sondrio, Italy) - an alpine ethnobotanical study. *J Ethnopharmacol.* 2013;145:517–29. <https://doi.org/10.1016/j.jep.2012.11.024> Elsevier.
90. Kayani S, Ahmad M, Zafar M, Sultana S, Khan MPZ, Ashraf MA, et al. Ethnobotanical uses of medicinal plants for respiratory disorders among the inhabitants of Gallies - Abbottabad, Northern Pakistan. *J Ethnopharmacol.* 2014;156:47–60. <https://doi.org/10.1016/j.jep.2014.08.005> Elsevier.
91. Krichen L, Audergon JM, Trifi-Farah N. Assessing the genetic diversity and population structure of Tunisian apricot germplasm. *Sci Hortic (Amsterdam).* 2014;172:86–100 Elsevier B.V. Available from: <http://www.sciencedirect.com/science/article/pii/S0304423814001769>.
92. Holdnack JA, Millis S, Larrabee GJ, Iverson GL. Assessing performance validity with the ACS. In: Holdnack JA, Drozdzik LW, Weiss LG, Iverson GL, editors. *WAIS-IV, WMS-IV, ACS Adv Clin Interpret;* Elsevier Inc; 2013. p. 331–65. <https://doi.org/10.1016/B978-0-12-386934-0.00007-9>.
93. Gagné F. Descriptive statistics and analysis in biochemical ecotoxicology. In: *Biochem Ecotoxicol Princ Methods;* Academic Press; 2014. p. 209–29.
94. Kichineni E, Wardle DA, Peltzer DA, Morse CW, Freschet GT. Contrasting effects of plant inter- and intraspecific variation on community-level trait measures along an environmental gradient. *Funct Ecol.* 2013;27:1254–61.
95. Tiwari SP, Kumar P, Yadav D, Chauhan DK. Comparative morphological, epidermal, and anatomical studies of *Pinus roxburghii* needles at different altitudes in the North-West Indian Himalayas. *Turk J Bot.* 2013;37:65–73.
96. Davis MB, Shaw RG. Range shifts and adaptive responses to quaternary climate change. *Science.* 2001;292:673–9.
97. Ingty T, Bawa KS. Climate change and indigenous people. In: Arrawatia ML, Tambe S, editors. *Clim Chang Sikk Patterns, Impacts Initiat: Information and Public Relations Department, Government of Sikkim;* 2012. p. 424. Available from: http://www.sikkimforest.gov.in/climate-change-in-sikkim/17-Chapter_ClimateChange_and_Indigenous_people.pdf.
98. Chen I-C, Hill JK, Ohlemüller R, Roy DB, Thomas CD. Rapid range shifts of species of climate warming. *Science.* 2011;333:1024–6.
99. Lulekal E, Asfaw Z, Kelbessa E, Van Damme P. Ethnomedicinal study of plants used for human ailments in Ankober District, North Shewa Zone, Amhara Region, Ethiopia. *J Ethnobiol Ethnomed.* 2013;9:63 Available from: <http://ethnobiomed.biomedcentral.com/articles/10.1186/1746-4269-9-63>.
100. Leto C, Tuttolomondo T, La Bella S, Licata M. Ethnobotanical study in the Madonie Regional Park (Central Sicily, Italy) - medicinal use of wild shrub and herbaceous plant species. *J Ethnopharmacol.* 2013;146:90–112. <https://doi.org/10.1016/j.jep.2012.11.042> Elsevier.
101. Aziz MA, Khan AH, Adnan M, Izatullah I. Traditional uses of medicinal plants reported by the indigenous communities and local herbal practitioners of Bajau Agency, Federally Administrated Tribal Areas, Pakistan. *J Ethnopharmacol.* 2017;198:268–81. Available from: <https://doi.org/10.1016/j.jep.2017.01.024>. Elsevier Ireland Ltd
102. Angmo K, Adhikari BS, Rawat GS. Changing aspects of traditional healthcare system in Western Ladakh, India. *J Ethnopharmacol.* 2012;143:621–630. Available from: <https://doi.org/10.1016/j.jep.2012.07.017>. Elsevier.
103. Soelberg J, Jäger AK. Comparative ethnobotany of the Wakhi agropastoralist and the Kyrgyz nomads of Afghanistan. *J Ethnobiol Ethnomed.* 2016;12:24.
104. Akhtar N, Akhtar S, Kazim S, Khan T. Ethnomedicinal study of important medicinal plants used for gynecological issues among rural women folk in district Gilgit, Pakistan. *Nat Sci.* 2016;14:30–4.
105. Sher H, Aldosari A, Ali A, de Boer HJ. Economic benefits of high value medicinal plants to Pakistani communities: an analysis of current practice and potential. *J Ethnobiol Ethnomed.* 2014;10:1–16 Available from: <https://ethnobiomed.biomedcentral.com/articles/10.1186/1746-4269-10-71>.
106. Khan SM, Page S, Ahmad H, Shaheen H, Ullah Z, Ahmad M, et al. Medicinal flora and ethnoecological knowledge in the Naran Valley, Western Himalaya, Pakistan. *J Ethnobiol Ethnomed.* 2013;9:4 Available from: <http://ethnobiomed.biomedcentral.com/articles/10.1186/1746-4269-9-4>.
107. Hussain S, Malik F, Khalid N, Abdul M, Riaz H. Alternative and traditional medicines systems in Pakistan: history, regulation, trends, usefulness, challenges, prospects and limitations. In: *A Compend Essays Altern Ther; 2012.* p. 89–108. Available from: <http://www.intechopen.com/books/a-compendium-of-essays-on-alternative-therapy/alternative-and-traditional-medicines-systems-in-pakistan-history-regulation-trends-usefulness-chall>.
108. Mukherjee PK, Harwanrah RK, Bahadur S, Banerjee S, Kar A, Chanda J, et al. Development of Ayurveda – tradition to trend. *J Ethnopharmacol.* 2017;197:10–24. <https://doi.org/10.1016/j.jep.2016.09.024>. Elsevier.
109. UNESCO. Culture and health orientation texts on the 1996 theme. In: *World Decad. Cult. Dev;* 1988–1997.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

