





Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Review

Trends in diabetes care with special emphasis to medicinal plants: Advancement and treatment

Bichitrananda Tripathy ^{a,*}, Nityananda Sahoo ^a, Sudhir Kumar Sahoo ^b^a Centurion University of Technology & Management, Odisha, 752050, India^b Royal College of Pharmacy and Health Sciences, Berhampur, Odisha, 759024, India

ARTICLE INFO

Keywords:

Diabetes mellitus
Plants
Pancreas
Insulin
Glucose

ABSTRACT

Diabetic mellitus (DM) is a common metabolic disorder prevailing throughout the world. It may affect a child to an older person depending upon the physiology and the factors influencing the internal metabolic system of the body. Several treatments are available in the market ranges from synthetic drugs, insulin therapy, herbal drugs, and transdermal patches. Interestingly, the development of technologies and digital health have proving very helpful in improving the lifestyle of diabetic patients. All treatment approaches have their own advantages and disadvantages in the form of effectiveness and side effects. Medicinal plants have a long history of traditional application in the treatment of diabetes and even the use of plants are growing day-by-day due to the significant results against diseases and fewer side effects as compared to other treatment therapies. The intention behind writing this review is to gather all information and discussed them exhaustively in an article. The novel Coronavirus 2019 (COVID-19) pandemic has affected my lives including diabetic patients. The antidiabetic treatment strategies during this period has also discussed. In this article, we highlighted the molecular mechanism and herbal phytoconstituents that are responsible for lowering blood glucose level. The factors responsible for the progression of metabolic disorders can be controlled with the use of phytoconstituents present in herbal plants to maintain β -cells performance and restore blood glucose level. It can be concluded that medicinal plants are effective and affordable with lesser side effects for treating DM.

1. Introduction

Diabetes mellitus (DM) has become a serious metabolic disorder and threatening public health all-over-the world. DM is among the top 10 deadly disorder in adults and estimated to cause four million deaths every year globally (Saeedi et al., 2019). According to the results of the International Diabetes Federation 2019, the global diabetes prevalence in 2019 was 463 million people. It is expected to rise up to 578 million by 2030 and 700 million by 2045 (Saeedi et al., 2019). DM is a condition of carbohydrate and lipid metabolism imbalance that increases blood glucose levels. Mechanistically, the body cannot regulate the amount of glucose in the blood during DM. It leads to hyperglycemia due to rise in carbohydrate contents (Peixoto Araujo et al., 2021). The two main types of diabetes are type 1 diabetes mellitus (T1DM), and type 2 diabetes mellitus (T2DM). T1DM is insulin-dependent DM generally occurs in children and youngsters; in this condition pancreas does not produce sufficient insulin. On the other hand, T2DM is non-insulin-dependent DM that occurs when the body cannot effectively use the

amiable insulin. T2DM has reached to epidemic proportions due to sedentary lifestyle and over-nutrition. It is very common in adults. It is considered as a complex disease that requires continuous medical care and strategies to reduce blood glucose level. Several biochemical and chemical agents are known that helps in controlling diabetes but no cure is yet available to get complete recover from the disorder (Verma et al., 2018). Education and support are very critical in controlling acute phases of diabetes and reducing risk of long-term complications (ADA, 2021). The treatment of diabetes has been improving day-by-day due to new research and technology that improving the well-being and health of diabetic patients. The field of diabetes care has been changing rapidly due to new research, technology, and treatments. These can improve the health and well-being of people with diabetes. The conventional treatments are no doubt effective but associated with certain complications or side effects. The complications are thus promoting researchers to develop some safe, affordable and efficient treatments against diabetes.

* Corresponding author.

E-mail address: bichitratripathy@gmail.com (B. Tripathy).

Interestingly, traditional herbal medicines have been found effective against diabetes (Kumar et al., 2017; Rathore et al., 2014). The phytoconstituents from extract have always been a rich source for controlling disorder and other complications arises due to it (Kumar et al., 2016). Medicinal plants have been used in the treatment of diabetes since ancient times (Singh et al., 2016). The World Health Organization (WHO) has found that 21,000 medicinal plants have been used for the treatment of different diseases all over the world, in which 2500 species of plants is found in India. The selection of herbs against diabetes depends on various factors such as stage of diabetes progression, availability, types of co-morbidities in patients, toxicities and safety profile of the herbs (Jain et al., 2014, 2015, 2016; Shrawan et al., 2015, 2016). The objective of the present review was to describe the utility of medicinal plants in the treatment of diabetes mechanistically. Thus, this article may help researchers in conducting future studies for the development of antidiabetic drugs and other therapies in the management of diabetes based on phytoconstituents and extracts from plants.

2. Factors causing diabetes

Diabetes is more prevailing in developing or low-income countries as compared to high-income nations, it is due to lifestyle factors and urbanization (IDF, 2017). The modifiable risk factors are obesity, hyperlipidemia, consumption of alcohol and tobacco (Barik et al., 2016). The vulnerable factors responsible for development of diabetes are biological risk factors including sex, age, marriage status, race, stress, and family history of diabetes; financial factors; severity of disease including poor disease control, complications, and co-morbidities; lack of education or low literacy; distrust of primary health services; occupational restriction; limited daily life behaviors; lifestyle factors including consumption of unhealthy food, lack of exercises and low-quality sleep; and mental condition including lack of friends and family and community environment (Chen et al., 2019; Bosun-Arije et al., 2019). Fig. 1 showing several genetic and environmental factors causing loss of β -cell mass and that leads to hyperglycemia. In this state, insulin do not respond to its demand in the body that causes increase in the level of blood glucose sufficient to diagnose diabetes (Skyler et al., 2017).

3. Pharmacologic therapy for treating diabetes

3.1. Treatment for type 1 diabetes mellitus (T1DM)

T1DM is a condition of serious decrease in the level on endogenous insulin and its release from β -cells of pancreas (Oram et al., 2019). The precise pathological mechanism and etiology are still not clear. It usually develops in child or young ones. People with type 1 diabetes are treated with multiple injection of prandial and basal insulin. The rapid acting insulin analogues are also used for reducing hypoglycemic risk. The rapid-acting prandial analogues i.e. lispro, aspart and glulisine are preferred over regular human insulin. However, aspart as faster-acting insulin is a new option due to advantage of better postprandial glucose coverage (Janež et al., 2020). The diabetic patients are being aware about prandial insulin dose to pre-meal blood glucose, carbohydrate intake and anticipated physical activity. The surgical treatment of type 1 diabetes such as islet and pancreas transplantation can also normalize the glucose levels and mitigate the complication of the disease (Meloche et al., 2007). However, patients availing this treatment need to take life long immunosuppressant drugs to prevent graft rejection of islet and pancreas.

3.2. Treatment for type 2 diabetes mellitus (T2DM)

T2DM is chronic metabolic disorder that develops due to defective insulin secretion and action. The primary cause of fasting hyperglycemia is the excessive production of glucose in the presence of hyperinsulinemia (ADA, 2013). However, after meals the impaired suppression of glucose production by insulin leading to decrease in the uptake of glucose by muscles and tissues. Metformin being the most preferred oral antidiabetic drug for the treatment of type 2 diabetes. It should be continued for life-long and not contraindicated with insulin. The second generation sulfonylureas include glyburide, glipizide and glimepiride; meglitinides include repaglinide, and nateglinide; thiazolidinediones include pioglitazone and rosiglitazone; alpha-glucosidase inhibitors include acarbose and miglitol; depeptidyl peptidase-4 inhibitors include sitagliptin, saxagliptin, linagliptin and alogliptin; colesevelam as bile acid sequestrant; bromocriptin as dopamine-2 agonist;




Fig. 1. Genetic and environmental factors causing hyperglycemia.

pramlintide as amylin mimetic; sodium glucose cotransporter - 2 inhibitors include canagliflozin, dapagliflozin, and empagliflozin; and glucagon-like peptide-1 receptor agonists include exenatide, liraglutide albiglutide and dulaglutide (Chamberlain et al., 2017). Table 1 showing the list of antidiabetic drugs with their mechanism and side effects. The injection of insulin is also considered during evidence of ongoing weight loss, symptoms of hyperglycemia (≥ 300 mg/dL). Fig. 2 showing pharmacological treatment for diabetes.

4. Medicinal plants used against diabetes traditionally

Medicinal plants have been used in the treatment of diseases since ancient times (McGaw et al., 2019). The importance of plants as raw material for pharmaceutical industries cannot be overlooked. The use

Table 1
List of existing anti-diabetic drugs with their mechanisms and side effects.

S. No.	Class	Mechanism of action	Side effects	Complication/Toxicities
1	Biguanide (Metformin)	Increases the effect of insulin	Reduced vitamin B12 absorption, weight loss, lactic acidosis, diarrhea, abdominal cramps	Liver failure, chronic kidney disease
2	Sulfonylureas (glyburide, glimepiride)	Enhances insulin secre- tion from pancreatic beta-cell ls	Hemolysis, agranulo- cytosis, weight gain	Obesity, cardiovascular comorbidities, allergy
3	Meglitinides (nateglinide, rep- aglinide)	Increase insulin secre- tion from pancreatic beta-cell ls	Weight gain, risk of hypoglycemia	Liver and renal failure
4	DPP-4 inhibitors (saxagliptin, sitagli- ptin)	Inhibit glucagon- like peptide- 1 degra- dation and promotes glucose-de- pendent insulin secretion	Pancreatitis, headache, gastrointestinal complaints, arthralgia, dizziness	Liver and renal failure
5	GLP-1 agonists (exenatide, liraglutide, albiglutide)	Direct stimulation of glucagon- like peptide- 1 receptor	Pancreatitis, pancreati c cancer, nausea	Gastrointestinal motility disorders
6	SGLT-2 inhibitors (canagliflozin, dapagliflozin, empagliflozin)	Increased glucosuria by inhibiting SGLT-2 in the kidney	Urinary tract infections, polyuria, dehydration, yeast infections, diabetic ketoacidosis	Urinary tract infections, chronic kidney disease
7	Alpha-glucosidase inhibitors (acarbose)	Reduce intestinal glucose absorption	Flatulence, diarrhea	Renal failure, inflammatory bowel disease
8	Thiazolidinediones (pioglitazone)	Reduce insulin resistance through the stimulation of PPARs	Cardiac failure, weight gain, osteoporosis, edema	Liver disease, congestive heart failure
9	Amylin analogues (pramlintide)	Reduce glucagon release	Risk of hypoglycemia, nausea	Gastroparesis

DPP-4, Dipeptidyl peptidase-4; SGLT-2, Sodium-glucose co-transporter-2; GLP-1, Glucagon-like peptide-1; PPARs, Peroxisome proliferator-activated receptors.

of medical plants is increasing worldwide due to its better long-term results with fewer side effects. They have been used to maintain and augment mental, spiritual and physical health along with treating diseased condition and ailments. Plants possess several active constituents present in their secondary metabolites such as alkaloids, phenol, glycosides, tannins, flavonoids, and terpenoids that are responsible for varieties of biological effects (Akinyemi et al., 2018). Thus, medicinal plants have an immense role to play in sustainable human health. Table 2 showing the list of medicinal plants having reported antidiabetic property.

4.1. *Acacia arabica*

Acacia arabica has been used traditionally in the treatment of diabetes, kidney and liver problems, healing wounds, and preventing of microbial attack. The mechanism of its antidiabetic action is due to presence of polyphenols in its bark. Its gives hypoglycemic effect by improving insulin sensitivity and reduction of fatty acid synthesis. Fisetinidol and robinetinidol are the two phytoconstituents responsible for antidiabetic action (Ikarashi et al., 2011).

4.2. *Bambusa arundinasia*

Bambusa arundinasia has been used traditionally for the treatment of diabetes, dysentery, peptic ulcer and aphthous. Mechanistically, it shows, α -glucosidase and α -amylase enzyme inhibitor action. Additionally, showing regeneration of pancreatic tissue and improvement in serum insulin level. It has also shown reduction of HbA1c (glycated hemoglobin), fructose-1-6-biphosphatase, glucose-6-phosphatase, triglycerides and total cholesterol. Stigmasterol and β -sitosterol are the two major compounds responsible for antidiabetic activity (Nazreen et al., 2011).

4.3. *Boswellia carterii*

Boswellia carterii is helpful in diabetes due to presence of oleo-gum resin. It shows antidiabetic activity by increasing liver glycogen, serum insulin and prevents degenerative change in β -cells of pancreas. *Boswellia carterii* shows hypoglycemic activity due to suppression of proinflammatory cytokine, apoptosis of cells, and lymphocytes penetration into pancreatic islets. The boswellic acid is the major activity phytoconstituent that showing inhibitor activity on diabetic complication following polyol aldose reductase pathway (Rao et al., 2013).

4.4. *Chrysanthemum indicum*

Chrysanthemum indicum has been used since a long term ago in the treatment of hypertension, inflammation and respiratory disorders in various countries. The phytoconstituents present in plant are luteolin and linarin that are responsible for anti-inflammatory effect on prostatitis and chronic pelvic inflammation. It is due to reduced activation of tissue necrosis factor - alpha (TNF- α) and interleukin-1beta (IL-1 β) (Dasgupta, 2019). *Chrysanthemum morifolium* Ramat. a species of *Chrysanthemum* used for antidiabetic activity (Yamamoto et al., 2015). It is a leading flowering plant with applied value worldwide (Su et al., 2019).

4.5. *Coriandrum sativum*

Coriandrum sativum is mainly used for its carminative property. However, its fruits have shown antidiabetic effect via enhancement of insulin release from β -cells along with increasing glucose uptake, glycogenesis and glucose oxidation. *C. sativum* improves cardio-protective indices and atherosclerotic index (Parseayan, 2012).




Fig. 2. Pharmacological treatment approaches for T1DM
SGLT-2, Sodium-glucose co-transporter-2.

4.6. *Glycyrrhiza glabra*

The antidiabetic activity of *G. glabra* is due to presence of glycyrrhetic acid present in roots of the plant. It increases expression of lipoprotein lipase in body tissues and improves sensitivity of insulin. It reduces total cholesterol, fatty acids in serum, and lipid deposition (Eu et al., 2010).

4.7. *Lactuca sativa*

The seeds of *L. sativa* has been used for its antidiabetic property by Persians since ancient times. It possesses alpha-glucosidase and alpha-amylase enzyme inhibitory activity. The mechanism of glucose lowering activity is based on elevation of serum insulin level, reduction of fructose-1-6- biphosphatase and glucose-6-phosphatase (Ahmed et al., 2013).

4.8. *Myrtus communis*

The leaves of *M. communis* possesses antidiabetic action by enhancing antioxidant function in liver of animals. The alpha-glucosidase enzyme inhibitory activity is the mechanism of its antidiabetic activity (Sepici-Dincel et al., 2007).

4.9. *Portulaca oleracea*

The seeds of *P. oleracea* shows hypoglycemic activity by reducing fasting and post-prandial blood glucose level. The mechanism of its antidiabetic activity is based on inhibition of alpha-glucosidase and alpha-amylase enzymes (Heidarzadeh et al., 2013).

4.10. *Punica granatum*

The flowers of *P. granatum* has been used for antidiabetic action, it is due to its alpha-glucosidase enzyme inhibitory activity and improvement of insulin sensitivity. It also enhances GLUT4 (Glucose Trans-

porter 4) and PPAR- γ (Peroxisome Proliferator-Activated Receptor- γ) (Huang et al., 2005).

4.11. *Rosa damascena*

The fruits and flowers of *R. damascena* possess glucose lowering and alpha-glucosidase enzyme inhibitory activity.

4.12. *Vitis vinifera*

V. vinifera shows antidiabetic activity due to presence of procyanidins in seeds. It also shows insulinomimetic activity via stimulation of insulin pathway mediator and upgrading glucose uptake. The pharmacological mechanism of plant against chronic diabetes include suppression of oxidative stress, reduction in advanced end glycation products (Li et al., 2008).

5. Highly valuable drug therapies and medicinal plants in 2020–21 for treating diabetes

The coronavirus disease 2019 (COVID-19) has imposed a global economy loss at various levels. During coronavirus disease 2019 (COVID-19) pandemic the diabetic patients were at increased risk of complication like Adult Respiratory Distress Syndrome and multi-organ failure. The following drugs were found more effective and having metabolically interfering effects against COVID-19 patients with diabetes: Metformin; Sodium-glucose-co-transporter-2 inhibitor including dapagliflozin, canagliflozin, and empagliflozin; Glucagon-like peptide-1 receptor agonists including dulaglutide, albiglutide, liraglutide, semaglutide, and lixisenatide; Dipeptidyl peptidase-4-inhibitors including linagliptin, alogliptin, sitagliptin, and saxagliptin; and Insulin (Bornstein et al., 2020). Vaccination should be prioritized in these patients. However, routine vaccination against Hepatitis B, influenza and pneumococcal pneumonia is recommended in diabetic patients (Pal et al., 2021). The current trends prevailing in the market under numerous segments as alternatives to injectable diabetes care. Based on product

Table 2

List of medicinal plants with reported antidiabetic activity.

S.No.	Plant	Family	Chemical constituents	Antidiabetic action	References
1	<i>Abies pindrow</i>	Pinaceae	Globulol, maltol, borneol, linalool	Stimulation of insulin secretion	Sinha et al., 2019
2	<i>Acacia arabica</i>	Fabaceae	Mallic acid, chlorogenic acid, catechin, epicatechin, chiorogenic acid, ellagic acid, corosolic acid,	Increasing the release of insulin	Hegazy et al. (2013), Bharti et al. (2018)
3	<i>Achyranthes rubrofusca</i>	Amaranthaceae	Betulinic acid, momordin Ib, zingibroside R1, Achyranthoside IV	Reduction of blood glucose level and restore pancreatic enzymatic activity	Geetha et al. (2011); He et al. (2017)
4	<i>Agrimonia eupatoria</i>	Rosaceae	4-caffeoylequinic acid, 3,5-dicaffeoyl quinic acid and luteolin-7-O-glucoside	Insulin-like and insulin releasing activity	Kuczmannová et al. (2016)
5	<i>Albizia lebbeck</i>	Fabaceae	Albiziazahexoside, lupeol, docosanoic acid, oleanolic acid, betasitosterol	α -amylase and α -glucosidase inhibitor activity	Verma et al. (2013); Ahmed et al. (2014)
6	<i>Aloe vera</i>	Asphodelaceae	lophenol, and cycloartanol	Inhibition of α -amylase and α -glucosidase enzymes	Tanaka et al. (2006); Muñiz-Ramirez et al. (2020)
7	<i>Amaranthus tricolor</i>	Amaranthaceae	Betacyanins, betaxanthins, rutin, isoquercetin, ferulic, ellagic, p-coumaric	Anti- α -amylase and anti- α -glucosidase activity	Yang et al. (2020); Peter and Gandhi (2017)
8	<i>Anacardium occidentale</i>	Anacardiaceae	Anacardic acid, cardanol, cardol	α -glucosidase inhibitor	Jaiswal et al. (2016); Palheta and Ferreira (2018)
9	<i>Annona squamosa</i>	Annonaceae	Acetogenin, annonacin, annonastatin samaquasine, squamone, acetogenin	Increase level of plasma insulin	Shirwaikar et al. (2004); Zahid et al. (2018)
10	<i>Averrhoa bilimbi</i>	Oxalidaceae	Cyanidin-3-o-h-glucoside, citric acids, amino acids	Increase level of plasma insulin	Kurup and S, 2017; Mathew et al. (2017)
11	<i>Azadirachta indica</i>	Meliaceae	Nimbidin	Normalizing altered insulin signaling molecules	Satyaranayana et al. (2015)
12	<i>Barleria prionitis</i>	Acanthaceae	Balarenone, pipataline, prioniside B, lupeol	Decrease blood glucose and increase serum insulin level	Dheer and Bhatnagar (2010); Shukla and Gunjegaonkar (2018); Banerjee et al. (2012)
13	<i>Bauhinia thonningii</i>	Fabaceae	Alepterolic acid, methyl-ent-3 β -hydroxylabd-8(17)-en-15-oate, 2 β -methoxyclovlan-9 α -ol (1)	Insulin mimetic activity	Salehi et al. (2019)
14	<i>Bixa orellana</i>	Bixaceae	Cryptoxanthin, phytoene, lutein, zeaxanthin	Promote binding of insulin on receptor and increase plasm insulin level	Vilar et al. (2014)
15	<i>Boerhaavia diffusa</i>	Nyctaginaceae	β -Sitosterol, arachidic acid, Ecdysone, urosilic acid, palmitic acid, hexacosonoic, β -triacontanol, hentriacontane	Increase level of plasma insulin	Alam et al. (2018)
16	<i>Caesalpinia ferrea</i>	Fabaceae	Galactomannan, kaempferol, galloylorientin, heptacosan	α -Amylase enzyme inhibitory action	Hassan et al. (2015); Macédo et al. (2020)
17	<i>Camellia sinensis</i>	Theaceae	Catechins, theaflavins, and caffeine	α -Amylase and α -Glucosidase inhibitory activity	Fu et al. (2017); Ardiana et al. (2017)
18	<i>Capsicum frutescens</i>	Solanaceae	Octadecadienal (Z), tetracosane, 3-carene, 5-eicosene, docosane	Stimulation of insulin secretion	Dougnon and Gbeassor (2016); Gurnani et al. (2016)
19	<i>Casearia esculenta</i>	Flacourtiaceae	3-hydroxymethyl xylitol	Insulin mimetic action	Govindasamy et al. (2011); Prakasam et al. (2005)
20	<i>Cassia fistula</i>	Fabaceae	β -sitosterol, lupeol, kaempferol, fistulin, rhein, leucopelargonidin	Increase insulin-stimulated glucose uptake	Einstein et al. (2013); Rahmani et al., 2015
21	<i>Cassia grandis</i>	Fabaceae	Linalool	α -glycosidase inhibitory activity	Loeha et al. (2010); Prada et al. (2018)
22	<i>Catharanthus roseus</i>	Apocynaceae	Limonene, dötriacontane, geraniol, citral, phytol	Increased secretion of insulin from β -cells pancreas	Rasineni et al. (2010); Lawal et al. (2015)
23	<i>Cecropia pachystachya</i>	Urticaceae	C-glycosyl flavonoids, proanthocyanidins	Increase secretion of insulin	Rivera-Mondragón et al. (2017); Costa, 2011
24	<i>Ceriops decandra</i>	Rhizophoraceae	Ceriopsis, lupeol, oleanolic acid, α -amyrin, ursolic acid, and catechin	Regeneration of β cells	Mahmud et al. (2018); Nabeel et al., 2010; Arora et al. (2014)
25	<i>Chiliadenus iphonoides</i>	Asteraceae	1,8-cineole, camphor, α -terpineol, terpin-4-ol, and bornyl formate	Increase secretion of insulin	Salehi et al. (2019); Abdelhalim et al., 2020
26	<i>Cinnamomum cassia</i>	Lauraceae	Cinnamaldehyde	Insulin mimetic action	Elumalai et al. (2011); Yan et al. (2015)
27	<i>Citrullus colocynthis</i>	Cucurbitaceae	Cucurbitacin L, isovitexin, khekadaengoside E	Restoration of pancreatic β -cells	Zheng et al. (2020); Dhakad et al. (2017); Shi et al. (2014)
28	<i>Clausena anisata</i>	Rutaceae	β -pinene, germacrene-D, sabinene, linalool, estragole	Stimulation of insulin secretion	Yakoob et al. (2016); Govindarajan, 2010
29	<i>Coscinium fenestratum</i>	Menispermaceae	Hentriacontan, palmitic acid, β -sitosterol, oleic acid	Decrease gluconeogenesis and increase enzymatic activity	Nayak et al. (2012); Malarvili et al. (2011)
30	<i>Eucalyptus citriodora</i>	Myrtaceae	Citronellol acetate, cis-geraniol, dihydrocarveol acetate, β -bisabolene, 3-hexen-1-ol, and pregn-5-en-20-one,3,17-dihydroxy-3-acetate	Increase glucose transporter 4 (GLUT-4) translocation	Wang et al. (2014); Dey and Mitra (2013)
31	<i>Eucalyptus globulus</i>	Myrtaceae	1,8-cineol, α -pinene	Enhance release of insulin from clonal pancreatic beta line	Chakraborty et al. (2018); Sebei et al. (2015)
32	<i>Ficus religiosa</i>	Moraceae	n-octacosanol, β -sitosteryl-D-glucoside, stigmasterol, lanosterol, methyl oleanolate, lupen-3-one	Stimulation of insulin secretion	Pandit et al. (2010); Chandrasekar et al. (2010)
33	<i>Gymnema sylvestre</i>	Apocynaceae	Gymnemic acid, gurmarin, tartaric acid, glucose, calcium oxalate, betaine, stigmasterol, and choline	Prevents absorption of glucose by the intestine to reduce blood sugar level	Kanetkar et al. (2007); Khan et al. (2019); Tiwari et al. (2014)

(continued on next page)

Table 2 (continued)

S.No.	Plant	Family	Chemical constituents	Antidiabetic action	References
35	<i>Heinsia crinata</i>	Rubiaceae	Sapogenin, neochlorogenin and diosgenin	Insulin level elevating effect	Okokon et al. (2009); Yobe et al. (2017)
36	<i>Helicteres isora</i>	Sterculiaceae	Kaempferol 7-O-coumaroylhexoside, rosmarinic acid and kaempferol 7-O-rhamnosylhexosides	Increase uptake of glucose	Olivas-Quintero et al. (2017)
37	<i>Hibiscus rosa</i>	Malvaceae	Beta-carotene, anthocyanin, Beta-sitosterol, arabinogalactans, gossypetin, L-ascorbic acid, citric acid	Stimulation of insulin secretion from beta pancreatic cells	Moqbel et al. (2011)
38	<i>Ipomoea batata</i>	Convolvulaceae	Caffeic acid, chlorogenic acid, rutin, quercetin	Decrease insulin resistance	Akhtar et al. (2018); Zengin et al. (2017)
39	<i>Juniperus communis</i>	Pinaceae	Longifolene, totarol, transcommunic acid	Stimulation of insulin secretion and increase glucose consumption	Raina et al. (2019)
40	<i>Momordica charantia</i>	Cucurbitaceae	Charantin, cucurbitacins, karounidiols, multiflorenol and nerolidol	Regulate glucose absorption by the gut and stimulate its uptake into muscles	Joseph and Jini (2013); Singh et al. (2011)
41	<i>Moringa oleifera</i>	Moringaceae	Oleic acid, ascorbic acid, 9-octadecenoic acid and 9-octadecenamide	α -glucosidase and pancreatic lipase inhibitory activity	Chen et al. (2020); Aja et al. (2014)
42	<i>Murraya koenigii</i>	Rutaceae	Linalool, geranyl acetate, elemol, allo-ocimene, myrcene, α -terpinene and neryl acetate	Increased the secretion of insulin and glycogenesis process	Rajendran et al. (2014); Sk et al. (2017)
43	<i>Olea europaea</i>	Oleaceae	Oleuropeoside	Increase uptake of glucose and release of insulin	Paramanick and Sharma (2017)
44	<i>Opuntia ficus-indica</i>	Cactaceae	Phytol, palmitate palmitic acid, and vitamin E	Increases glucose uptake through activation of AMPK/p38 MAPK pathway	Halmi et al. (2012); Luo et al. (2010); Leem et al. (2016)
45	<i>Origanum vulgare</i>	Lamiaceae	Amburosides A, apigenin 7-O-glucuronide, luteolin 7-O-glucuronide, lithospermic acid, rosmarinic acid, and demethylbenzolignanoid	α -glucosidase inhibitory activity	Yu et al. (2021);
46	<i>Passiflora nitida</i>	Passifloraceae	Luteolin, apigenin, kaempferol and quercetin	α -glucosidase inhibitory activity	Casierra-Posada and Jarma-Orozco (2016)
47	<i>Paspalum scrobiculatum</i>	Poaceae	Vanillic acid, syringic acid, cis- ferulic acid, p-hydroxy benzoic acid and melilotic acid	Increase glycogen synthesis and decrease in glycated haemoglobin levels	Kiran et al. (2014); Jain et al. (2009)
48	<i>Persea americana</i>	Lauraceae	Peptone, glycosylated abscisic acid, cellulose, b-galactoside, polyuronoids, and polygalacto urease	Inhibition of insulinase activity	Yasir et al. (2010); Ezejiofor et al. (2013)
49	<i>Phoenix dactylifera</i>	Arecaceae	β -carotene, ascorbic acid, α -tocopherols, selenium	Glucose lowering effect	Abdelaziz et al. (2015)
50	<i>Phyllanthus niruri</i>	Euphorbiaceae	Phyllanthin, coumarins, chlorogenic acids, and anthocyanins	Inhibition of glucose absorption and enhancement of glucose storage	Sibiya et al. (2020);
51	<i>Phyllanthus simplex</i>	Euphorbiaceae	Corilagin, gallic acid, phyllanthin, geraniin and niranthin	Glucose lowering effect	Mao et al. (2016)
52	<i>Picralima nitida</i>	Magnoliopsida	Akuammamine, akuammidine, akuammicine, pseudo-akuammagine	Regeneration of β cells	Teugwa et al. (2013);
53	<i>Piper longum</i>	Piperaceae	Piplartine and piperine	Glucose lowering effect	Singh and Navneet, 2018; Nabi et al. (2013)
54	<i>Scoparia dulcis</i>	Scrophulariaceae	scoparic acid D	Secretagogue activity of insulin	Latha et al. (2009)
55	<i>Sonchus oleraceus</i>	Asteraceae	Apigenin, luteolin, 1-cerotol, germanicyl acetate, and oleanolic acid	α -amylase and α -glucosidase inhibitory activity	Xu and Liang (2005); Li and Yang (2018)
56	<i>Swertia chirayata</i>	Gentianaceae	Amarogentin, gentianine, ursolic acid, isobellidifolin, sweroside, magniferin	Stimulation of insulin secretion from pancreatic islets	Dey et al. (2020)
57	<i>Syzygium jambolana</i>	Myrtaceae	Glucoside, anthocyanins, isoquercetin, kaemferol, ellagic acid and myrecetin	Stimulates insulin secretion	Ayyanar and Subash-Babu (2012)
58	<i>Tamarindus indica</i>	Fabaceae	Citric acid, tartaric acid, malic acid, acetic acid, formic acid, and succinic acid	α -amylase and α -glucosidase inhibitory activity	Bhadoriya et al. (2011); Krishna et al. (2020)
59	<i>Terminalia chebula</i>	Combretaceae	Chebulic acid, neo-chebulic acid mannitol, chebulagic acid, corilagin	Insulin mimetic action	Senthilkumar et al. (2006); Chang and Lin (2012)
60	<i>Terminalia catappa</i>	Combretaceae	Asiatic acid, vitexin, ursolic acid, isovitexin, gallic acid, tergalagin, tercatatin, punicalagin, chebulagic acid, punicalin, terflavins A and B, and geranin	α -glucosidase and α -amylase inhibitory activity	Mininel et al. (2014); Behl and Kotwani (2017)
61	<i>Tinospora crispa</i>	Menispermaceae	apeginin, diosmetin, genkwanin, cycloecalenol, cycloecalenone	Stimulation of insulin secretion from pancreatic islets	Klangjareonchai et al. (2012)
62	<i>Trigonella foenum-graecum</i>	Fabaceae	Diosgenin, trigonelline, gentianine, carpaine, butanoic acid, and isovaleric acid	Restoration of pancreatic β -cells	Wani and Kumar, 2018; Geberemeskel et al. (2019)
63	<i>Urtica dioica</i>	Urticaceae	Histamine, acetylcholine, 5-hydroxytryptamine	Stimulation of insulin secretion	El Haouari et al. (2019); Joshi et al. (2019)
64	<i>Vaccinium arctostaphylos</i>	Ericaceae	Linalool, α -Pinene, Safranal and Sandaracopimaradiene	Insulin level elevating effect	Salehi et al. (2019); Teimouri et al. (2015); Kianbakht and Hajighaee (2013)
65	<i>Vernonia amygdalina</i>	Asteraceae	Vernomygdin, Vernoniosides Vernodalol Vernodalin, Epivernodalol	Suppression of gluconeogenesis	Atangwho et al. (2009); Atangwho et al. (2014)
66	<i>Vinca rosea</i>	Apocynaceae	Vincristine, vinblastine	Regeneration and rejuvenation of beta cells	Ahmed et al. (2010)

(continued on next page)

Table 2 (continued)

S.No.	Plant	Family	Chemical constituents	Antidiabetic action	References
67	<i>Zaleya decandra</i>	Aizoaceae	6-octadecenoic acid, n-hexadecanoic acid	Stimulation of insulin secretion	Meenakshi et al. (2010)
68	<i>Zingiber officinale</i>	Zingiberaceae	Gingerols, paradols, shogaols, gingerdiones, zingerones, gingerdiols	Stimulation of insulin secretion and decrease glucose level	Obih et al. (2017)
69	<i>Zizyphus mauritiana</i>	Rhamnaceae	Palmitic acid, ethyl stearate and α -linolenic acid	Restoration of blood glucose level	Ashraf et al. (2015); Niamat et al. (2012)

GLUT-4, Glucose transporter-4; AMPK, AMP-activated protein kinase; MAPK, Mitogen-activated protein kinase.

types, it include insulin pumps, smart glucose meter, tethered insulin pumps, implantable insulin pumps, insulin patches and artificial pancreas. Based on device types include wearable and hand held.

Several plants have been used for treating diabetes and continue to get most beneficial agents with lesser side effects. Interestingly, few plants are very common in different countries that are frequently used in diabetes due to their effectiveness, economic profile and easy availability. The passed year was very difficult due to COVID pandemic throughout the world; peoples stayed at their homes and protecting themselves. However, medicines were accessible to patients even though the diabetic patients have been consuming herbs to remain healthy and maintain normal blood glucose levels. Famous medicinal plants used for antidiabetic purposes in the year 2020–21 are as follows:

5.1. Aloe vera

Aloe vera L. has been used traditionally in the treatment of various diseases in many countries since a long term ago. The major chemical constituents present in the plant are anthraquinones, phytosterols, lignins, salicylic acid and polysaccharides. It is very efficient in reducing blood glucose level of diabetic patients by improving the body response towards insulin. *A. vera* is also effective in reducing lipid level in the body due to presence of phytosterols such as lupeol, campesterol and sitosterol that are structurally similar to cholesterol and reduce its absorption by competition with it (Joseph et al., 2018; Choudhary et al., 2014).

5.2. Cinnamon

Cinnamomum cassia is commonly known as cinnamon. The bark of cinnamon is sweet in taste and most widely used as flavoring agents in the food and beverages. It is well recognized due to its medicinal properties and traditionally used in menstrual irregularities, diarrhea and arthritis. Cinnamon is a well known plant, its bark is used for antidiabetic action. Around 250 species of Cinnamon has reported in which *C. cassia* and *C. zeylanicum* are popular ones (Medagama, 2015).

5.3. Ginger

Ginger (*Zingiber officinale*) is a safe and non-toxic spice with lesser side effects as considered by the food and drug administration (Huang et al., 2019). Its origin is from Southeast Asia and has a long traditional use due to its medicinal properties. Ginger is used as an antidiabetic herb due to its action on inhibiting the enzymes α -glucosidase and α -amylase. It is due to the presence of shagol and gingerol as phytoconstituents (Lindstedt, 2018).

5.4. Garlic

Garlic (*Allium sativum*) belongs to family Liliaceae. *A. sativum* is used exhaustively due to its medicinal and nutritional properties. It is native to central Asia and being used for culinary purposes due to its health benefits since ancient times. The phytoconstituents present in plant parts are Diallyl disulfide, Allicin, S-allyl cysteine, Allyl mercaptan and

Alliin. *A. sativum* is effective in reducing blood glucose level and insulin resistance in the body. The other well established uses of garlic include anticoagulant, antiatherosclerotic, anticancer, antimicrobial, immunomodulatory, antidote and hypolipidemic activities (Shabani et al., 2019).

5.5. Bitter gourd

Bitter gourd is *Momordica charantia* Linn. belongs to family Cucurbitaceae. It is also known as karela and balsam pear. The fruits of *M. charantia* has been used for several decades to treat diabetes to the indigenous people of Asia, East Africa and South America. The hypoglycemic mechanism of *M. charantia* is due to presence of phytoconstituents that possess AMP-activated protein kinase activity. The phytoconstituents present in the plant are charantin, cucurbitins, karounidols, momordicinin, stigmasterol, zeaxanthin and pipecolic acid (Joseph and Jini, 2013; Mahmoud et al., 2017).

5.6. Sweet potato

Sweet potato (*Ipomoea batatas*) is used as an edible product in various countries including India, Tanzania, Japan, Peru and New Zealand. The shoots and leaves have been employed traditionally as medicine in the treatment of diabetes. It belongs to the family Convolvulaceae. The roots of *I. batatas* are rich in mineral salts and vitamin C. It is used to alleviate vitamin A deficiency in Eastern India and East Africa due to high quantity of beta-carotene in roots (Gunn et al., 2013; Kusano and Abe, 2000).

5.7. Black seed

Black seed (*Nigella sativa*) or black cumin is dicotyledon plant belongs to family Ranunculaceae. It has been used as an important culinary herb containing medicinal properties. The phytoconstituents present in the seeds are p-cymene, nigellone, flavonoids, carvone and thymoquinone in which thymoquinone is considered the chief active compound. The seeds are used as good antidiabetic agents that can lower increased glucose level of the body and also restore lipid profile. It is also used for neuroprotective, nephroprotective, anticonvulsant, anticancer and antimutagenic properties (Heshmati and Namazi, 2015; Bamosa et al., 2010; Gunn et al., 2013).

5.8. Chirayita

Chirayita (*Swertia chirayita*) is a popular indigenous herb of Himalayas belongs to family Gentianaceae. It has been traditionally used in the treatment of several ailment including diabetes, malaria and liver disorders. It is a rich sources of flavonoids, xanthones, terpenoids, iridoid and alkaloids. *S. chirayita* is known for its bitter taste, it is due to the presence of phytoconstituents such as amarogentin, swertiamarin, and swerchirin (Alam et al., 2011; Kumar and Van Staden, 2016; Dey et al., 2020).

6. Advancement in the treatment of diabetes

- An artificial pancreatic system is a big addition in the treatment of diabetes. It is critical to quantify blood glucose level non-enzymatically, accurately and in stable manner (Wang et al., 2019).
- Tepelizumab, an immune stimulant drug is under phase 3 trial that delayed type 1 diabetes. Anti-thymocyte globulin (ATG) is good alternative for people with newly diagnosed type 1 diabetes. It can preserve the function of beta cells and decreases blood sugar levels.
- A new protein, Hybrid Insulin Peptides (HIPS) has found on beta-cells of pancreas with type 1 diabetes and this are recognized as foreign body by the immune cells of patients (Baker et al., 2019). Thus, HIPS can be an important target for type 1 diabetic patients.
- Bisphenol A (BPA), a synthetic chemical present in our food has identified recently its association with increased risk for developing type 2 diabetes. Therefore, it should be administered in a controlled manner to humans due to its direct effect on glucose and insulin levels (Hagopian et al., 2019).
- The digital diabetes care market has gone beyond the earlier diabetes management. The digital solutions for diabetes such as applications, connected devices and services along with healthcare professionals and payers have change the earlier scenario of the disease (Deepa et al., 2020). Digital health is proving very helpful in improving the lifestyle of diabetic patients (Kaufman and Khurana, 2016). It is a complementary tool in diabetes-intervention studies.

7. Scope and prospects

Diabetes is a common metabolic disorder that can affect people of any group in developed as well as developing countries (Zaid et al., 2016). The rise in global diabetic patients has been linked with several factors like poor diet, obesity and sedentary lifestyles. However, older people have share a significant portion of diabetic patient worldwide. Several conventional and pharmacological treatments are available to restore normal blood glucose level. However, the conventional therapies and drugs treatments are associated with certain side effects. The advancement in the technologies like mobile phones and personal digital assistants are very helpful to tell the current status of health of an individual. The most frequently used digital services for healthcare wellness including mobile telemedicine, health call centers and emergency toll-free telephone services according to the second global survey on electronic health (WHO, 2011). However, a long-term study is desired to determine their sustainability, efficacy, cost-effectiveness and patient satisfaction. The traditional herbal medicines have been used in the treatment of several disorders (Jain et al., 2015; Rao et al., 2014, 2015; Rathore et al., 2014). The bioactive compounds are very effective in controlling bacteria, pathogens, viruses, and other foreign agents (Jain et al., 2019 a; Jain et al., 2018; Jain et al., 2020 a; Jain et al., 2020 b; Jain et al., 2020 c; Rao et al., 2016; Jain et al., 2020 d; Jain et al., 2020 e). The medicinal plants have been used in the treatment of diabetes and they are getting recognition after their standardization and proper clinical trials (Jain et al., 2019 b). The herbs have several modes of antidiabetic action including insulin mimetic, alteration of glucose metabolizing enzymes, inhibition of intestinal absorption of glucose, increase uptake of glucose peripherally, regeneration of pancreatic cells, promotes insulin release and ameliorating oxidative stress (Mwiti and Jide, 2015). Several herbs have been used traditionally in the treatment of diabetes since ancient times in many countries. It is due to their effective role in medicinal field. The herbal drugs are recognizing worldwide because of their safety profile improved pharmacokinetic, pharmacological and clinical status (Choudhury et al., 2017).

8. Conclusion

Medicinal plants have been used in the treatment of several disorders. They can lower the blood glucose level through combination of more than one mechanism. These plants have similar mode of action and mechanism as conventional drugs for treating diabetes but advantageously they have lesser side effects. There is a need to standardize all the traditional herbs used for medicinal purposes and determine their molecular mechanism for alleviating diabetes. The standardized, validated and characterized herbal drugs with their identified phytoconstituents can enter the world of clinical trials and further reach to their ultimate market. Thus, traditional knowledge for the use of medicinal herbs against diabetes with molecular mechanism may be a useful tool for new drug development.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgement

The authors are thankful for providing all facilities required for preparing the review article. We are showing our explicitly gratitude towards Probecell: Scientific Writing Services for proofreading and editing of this article.

References

- Abdelaziz, D.H.A., Ali, S.A., Mostafa, M.M.A., 2015. Phoenix dactylifera seeds ameliorate early diabetic complications in streptozotocin-induced diabetic rats. *Pharm. Biol.* 53 (6), 792–799.
- Abdelhalim, A., Al-Munawarah, A.-M., 2020. Pharmacological properties and chemical constituents of chilaidenus iphionoides (syn. Varthemia iphionoides): a review. *Eur. J. Med. Plants* 31, 84–97.
- ADA. American Diabetes Association, 2013. Diagnosis and classification of diabetes mellitus. *Diabetes Care* 36 (1), 67–74.
- ADA, 2021. American diabetes association. Standards of medical care in diabetes—2021. *Diabetes Care* 44 (1), 1–2.
- Ahmed, D., Kumar, V., Sharma, M., Verma, A., 2014. Target guided isolation, in-vitro antidiabetic, antioxidant activity and molecular docking studies of some flavonoids from Albizzia Lebbeck Benth. bark. *BMC Compl. Alternative Med.* 14 (1), 155.
- Ahmed, D., Sharma, M., Mukerjee, A., Ramteke, P.W., Kumar, V., 2013. Improved glycemic control, pancreas protective and hepatoprotective effect by traditional poly-herbal formulation “Qurs Tabasheer” in streptozotocin induced diabetic rats. *BMC Complement Altern Med.* 13, 10.
- Ahmed, M.F., Kazim, S.M., Ghori, S.S., Mehjabeen, S.S., Ahmed, S.R., Ali, S.M., et al., 2010. Antidiabetic activity of Vinca rosea extracts in alloxan-induced diabetic rats. In: Dandona, P.K. (Ed.), *Internet J. Endocrinol.* 2010, 841090.
- Aja, P.M., Nwachukwu, N., Ibiam, U.A., Igwenyi, I.O., Offor, C.E., Orji, U.O., 2014. Chemical constituents of moringa oleifera leaves and seeds from Abakaliki , Nigeria. *AJPCT* 2 (3), 310–321.
- Akhtar, N., Akram, M., Daniyal, M., Ahmad, S., 2018. Evaluation of antidiabetic activity of Ipomoea batatas L. extract in alloxan-induced diabetic rats. *Int. J. Immunopathol. Pharmacol.* 32 2058738418814678.
- Akinyemi, O., So, O., Ka, J., 2018. Medicinal plants and sustainable human health : a review. 2 (4), 194–195.
- Alam, K., Ali, S., Mahjabeen, S., Hassan, R., Rahman, M.F., Chowdhury, R., 2011. Potential hypoglycemic effect of Swertia chirayita—an Indian subcontinent herb with important medicinal value. *Pharmacologyonline* 2, 642–647.
- Alam, P., Shahzad, N., Gupta, A.K., Mahfuz, A.M., Bamagous, G.A., Al-Ghamdi, S.S., et al., 2018. Anti-diabetic effect of boerhavia diffusa L. Root extract via free radical scavenging and antioxidant mechanism. *Toxicol Environ Health Sci* 10 (3), 220–227.
- Ardiana, L., Sauriasari, R., Elya, B., 2017. Antidiabetic activity studies of white tea (*camellia sinensis* (L.) O. Kuntze) ethanolic extracts in streptozotocin-nicotinamide induced diabetic rats. *Pharmacogn J* 10, 186–189.
- Arora, K., Nagpal, M., Jain, U., Jat, R.C., Jain, S., 2014. Mangroves : a novel gregarious phyto medicine for diabetes. *Int. J. Res. Dev. Pharm. Life Sci.* 3 (6), 1231–1244.
- Ashraf, A., Sarfraz, R.A., Anwar, F., Shahid, S.A.L.I., Alkhafry, K.M., 2015. Chemical composition and biological activities of leaves of ziziphus mauritiana l . native to Pakistan. *Pakistan J. Bot.* 47 (1), 367–376.
- Atangwho, I., Ebong, P., Eyoung, E., Williams, I., Egbung, E., 2009. Comparative chemical composition of leaves of some antidiabetic medicinal plants: Azadirachta indica, Vernonia amygdalina and Gongronema latifolium. *Afr. J. Biotechnol.* 8.
- Atangwho, I.J., Yin, K.B., Umar, M.I., Ahmad, M., Asmawi, M.Z., 2014. Vernonia amygdalina simultaneously suppresses gluconeogenesis and potentiates glucose

- oxidation via the pentose phosphate pathway in streptozotocin-induced diabetic rats. *BMC Compl. Alternative Med.* 14, 426.
- Ayyanar, M., Subash-Babu, P., 2012. *Syzygium cumini* (L.) Skeels: a review of its phytochemical constituents and traditional uses. *Asian Pac J Trop Biomed* 2 (3), 240–246.
- Baker, R.L., Rihanek, M., Hohenstein, A.C., Nakayama, M., Michels, A., Gottlieb, P.A., Haskins, K., Delong, T., 2019. Hybrid insulin peptides are autoantigens in type 1 diabetes. *Diabetes* 68 (9), 1830–1840.
- Bamosa, A.O., Kaatabi, H., Lebdaa, F.M., Elq, A.M., Al-Sultanh, A., 2010. Effect of *Nigella sativa* seeds on the glycemic control of patients with type 2 diabetes mellitus. *Indian J. Physiol. Pharmacol.* 54 (4), 344–354.
- Banerjee, D., Maji, A.K., Mahapatra, S., Banerji, P., 2012. *Barleria prionitis* linn.: a review of its traditional uses, phytochemistry, pharmacology and toxicity. *Res. J. Phytochem.* 6, 31–41.
- Barik, A., Mazumdar, S., Chowdhury, A., Rai, R.K., 2016. Physiological and behavioral risk factors of type 2 diabetes mellitus in rural India. *BMJ Open Diabetes Res & Care* 4 (1), e000255.
- Behl, T., Kotwani, A., 2017. Proposed mechanisms of *Terminalia catappa* in hyperglycaemia and associated diabetic complications. *J. Pharm. Pharmacol.* 69 (2), 123–134.
- Bhadoriya, S.S., Ganeshpurkar, A., Narwaria, J., Rai, G., Jain, A.P., 2011. *Tamarindus indica*: extent of explored potential. *Pharmacogn Rev* 5 (9), 73–81.
- Bharti, S.K., Krishnan, S., Kumar, A., Kumar, A., 2018. Antidiabetic phytoconstituents and their mode of action on metabolic pathways. *Ther Adv Endocrinol Metab* 9 (3), 81–100.
- Bornstein, S.R., Rubino, F., Khunti, K., Mingrone, G., Hopkins, D., Birkenfeld, A.L., et al., 2020. Practical recommendations for the management of diabetes in patients with COVID-19. *Lancet Diabetes Endocrinol* 8 (6), 546–550.
- Bosun-Arije, F.S., Ling, J., Graham, Y., Hayes, C., 2019. A systematic review of factors influencing Type 2 Diabetes Mellitus management in Nigerian public hospitals. *Int J Africa Nurs Sci [Internet]* 11, 100151.
- Casierra-Posada, F., Jarma-Orozco, A., 2016. Nutritional composition of passiflora species. pp. 517–534.
- Chakraborty, M., Bagchi, B., Das, S., Basu, R., Nandy, P., 2018. A dose Dependent hepatoprotective and nephroprotective activity of eucalyptus oil on Streptozotocin induced diabetic mice model. *Clin Phytoscience* 4 (1), 10.
- Chamberlain, J.J., Herman, W.H., Leal, S., Rhinehart, A.S., Shubrook, J.H., Skolnik, N., et al., 2017. Pharmacologic therapy for type 2 diabetes: synopsis of the 2017 American diabetes association standards of medical care in diabetes. *Ann. Intern. Med.* 166 (8), 572–578.
- Chandrasekar, S.B., Bhanumathy, M., Pawar, A.T., Somasundaram, T., 2010. Phytopharmacology of *Ficus religiosa*. *Pharmacogn Rev* 4 (8), 195–199.
- Chang, C.L., Lin, C.S., 2012. Phytochemical composition, antioxidant activity, and neuroprotective effect of *Terminalia chebula retzii* extracts. In: Zhang, B. (Ed.), *Evidence-Based Complement Altern Med.* p. 125247.
- Chen, G.L., Xu, Y.B., Wu, J.L., Li, N., Guo, M.Q., 2020. Hypoglycemic and hypolipidemic effects of *Moringa oleifera* leaves and their functional chemical constituents. *Food Chem.* 333, 127478.
- Chen, J., Jing, X., Liu, X., Volkmann, A.-M., Chen, Y., Liu, Y., et al., 2019. Assessment of factors affecting diabetes management in the City Changing Diabetes (CCD) study in Tianjin. *Plos One* 14 (2), e0209222.
- Choudhary, M., Kochhar, A., Sangha, J., 2014. Hypoglycemic and hypolipidemic effect of *Aloe vera* L. in non-insulin dependent diabetics. *J. Food Sci. Technol.* 51 (1), 90–96.
- Choudhury, H., Pandey, M., Hua, C.K., Mun, C.S., Jing, J.K., Kong, L., et al., 2017. An update on natural compounds in the remedy of diabetes mellitus: a systematic review. *J Tradit Complement Med* 8 (3), 361–376.
- Deepa, M., Shruti, M., Mohan, V., 2020. Reducing the global burden of diabetes using mobile health. *Diabetes Digital Health. INC*, pp. 1–24.
- Dey, B., Mitra, A., 2013. Chemo-profiling of eucalyptus and study of its hypoglycemic potential. *World J. Diabetes* 4 (5), 170–176.
- Dey, P., Singh, J., Suluvoy, J.K., Dilip, K.J., Nayak, J., 2020. Utilization of swertia chirayita plant extracts for management of diabetes and associated disorders: present status, future prospects and limitations. *Nat Products Bioprospect* 10 (6), 431–443.
- Dhakad, P.K., Sharma, P.K., Kumar, S., 2017. A review on phytochemical studies and biological potential of *Citrullus colocynthis* (L.) schrad. (cucurbitaceae). *J* 5 (4), 55–64.
- Dheer, R., Bhatnagar, P., 2010. A study of the antidiabetic activity of *Barleria prionitis* Linn. *Indian J. Pharmacol.* 42, 70–73.
- Dougnon, T.J., Gbeassor, M., 2016. Evaluation of the effects of the powder of *Capsicum frutescens* on glycemia in growing rabbits. *Vet. World* 9 (3), 281–286.
- Einstein, J.W., Mohd Rais, M., Mohd, M.A., 2013. Comparative evaluation of the antidiabetic effects of different parts of *Cassia fistula* linn, a Southeast asian plant. In: Valention, P. (Ed.), *J. Chem.* 2013, 714063.
- El Haouari, M., Rosado, J.A., 2019. Phytochemical, anti-diabetic and cardiovascular properties of *urtica dioica* L. (urticaceae): a review. *Mini Rev. Med. Chem.* 19 (1), 63–71.
- Elumalai, S., Kesavan, R., Selvarajan, R., Murugasen, R., Elumalai, E., 2011. Isolation, purification and identification of the anti-diabetic components from *Cinnamomum zeylanicum* and *Cinnamomum cassia* bark oil extracts. *Curr. Bot. 2.*
- Eu, C.H., Lim, W.Y., Ton, S.H., et al., 2010. Glycyrrhetic acid improved lipoprotein lipase expression, insulin sensitivity, serum lipid and lipid deposition in high-fat diet-induced obese rats. *Lipids Health Dis.* 9, 81.
- Ezejiofor, A.N., Okorie, A., Orisakwe, O.E., 2013. Hypoglycaemic and tissue-protective effects of the aqueous extract of *persea americana* seeds on alloxan-induced albino rats. *Malays. J. Med. Sci.* 20 (5), 31–39.
- Fu, Q.Y., Li, Q.S., Lin, X.M., Qiao, R.Y., Yang, R., Li, X.M., et al., 2017. Antidiabetic effects of tea. *Molecules* 22 (5), 849.
- Geberemeskel, G.A., Debebe, Y.G., Nguse, N.A., 2019. Antidiabetic effect of fenugreek seed powder solution (*Trigonella foenum-graecum* L.) on hyperlipidemia in diabetic patients. In: Portha, B. (Ed.), *J Diabetes Res* 2019, 8507453.
- Geetha, G., Kalavalasarasi Gopinathapillai, P., Sankar, V., 2011. Anti diabetic effect of Achyranthes rubrofusca leaf extracts on alloxan induced diabetic rats. *Pak. J. Pharm. Sci.* 24 (2), 193–199.
- Govindarajan, M., 2010. Chemical composition and larvicidal activity of leaf essential oil from *Clausena anisata* (Willd.) Hook. f. ex Benth (Rutaceae) against three mosquito species. *Asian Pac J Trop Med* 3 (11), 874–877.
- Govindasamy, C., Al-Numair, K.S., Alsaif, M.A., Viswanathan, K.P., 2011. Influence of 3-hydroxymethyl xylitol, a novel antidiabetic compound isolated from *Casearia esculenta* (Roxb.) root, on glycoprotein components in streptozotocin-diabetic rats. *J. Asian Nat. Prod. Res.* 13 (8), 700–706.
- Gunn, J., Che, C.-T., Farnsworth, N., 2013. Chapter 33 - diabetes and natural products. In: Watson RR, Preedy VRBT-BF as DI for D. Academic Press, San Diego, pp. 381–394.
- Gurnani, N., Gupta, M., Mehta, D., Mehta, B.K., 2016. Chemical composition, total phenolic and flavonoid contents, and in vitro antimicrobial and antioxidant activities of crude extracts from red chilli seeds (*Capsicum frutescens* L.). *J Taibah Univ Sci* 10 (4), 462–470.
- Hagopian, T.A., Bird, A., Stanelle, S., Williams, D., Schaffner, A., Phelan, S., 2019. Pilot study on the effect of orally administered Bisphenol A on glucose and insulin response in nonobese adults. *Journal of the Endocrine Society* 3 (3), 643–654.
- Halmi, S., Benlakssira, B., Bechtarzi, K., Djerrou, Z., Djeaalab, H., Riachi, F., et al., 2012. Antihyperglycemic activity of prickly pear (*Opuntia ficus-indica*) aqueous extract. *Int. J. Med. Aromatic Plants* 2, 2249–2340.
- Hassan, S., El-Sammad, N., Moussa, A., Mohammed, M., Farrag, A.R., Hashim, A., et al., 2015. Hypoglycemic and Antioxidant Activities of *Caesalpinia Ferrea Martius Leaf Extract* in Streptozotocin-Induced Diabetic Rats.
- He, X., Wang, X., Fang, J., Chang, Y., Ning, N., Guo, H., et al., 2017. The genus Achyranthes: a review on traditional uses, phytochemistry, and pharmacological activities. *J. Ethnopharmacol.* 203, 260–278.
- Hegazy, G.A., Alnoury, A.M., Gad, H.G., 2013. The role of *Acacia Arabica* extract as an antidiabetic, antihyperlipidemic, and antioxidant in streptozotocin-induced diabetic rats. *Saudi Med. J.* 34 (7), 727–733.
- Heidarzadeh, S., Farzanegi, P., Azarbayjani, M.A., et al., 2013. Purslane effect on GLP-1 and GLP-1 receptor in type 2 diabetes. *Electron. Physician* 5, 582–587.
- Heshmati, J., Namazi, N., 2015. Effects of black seed (*Nigella sativa*) on metabolic parameters in diabetes mellitus: a systematic review. *Compl. Ther. Med.* 23 (2), 275–282.
- Huang, F., Deng, T., Meng, L., Ma, X., 2019. Dietary ginger as a traditional therapy for blood sugar control in patients with type 2 diabetes mellitus: a systematic review and meta-analysis. *Medicine (Baltimore)* 98 (13).
- Huang, T.H.W., Peng, G., Kota, B.P., et al., 2005. Anti-diabetic action of *Punica granatum* flower extract: activation of PPAR-g and identification of an active component. *Toxicol. Appl. Pharmacol.* 207, 160–169.
- IDF; International Diabetes Federation, 2017. *Diabetes Atlas*. eighth ed. IDF.
- Ikarashi, N., Toda, T., Okaniwa, T., Ito, K., Ochiai, W., Sugiyama, K., 2011. Anti-obesity and anti-diabetic effects of acacia polyphenol in obese diabetic KKAY mice fed high-fat diet. *Evide Based Complement Alternat Med* 2011, 952031.
- Jain, A., Jain, P., Parihar, D.K., 2019 b b. Comparative study of in-vitro antidiabetic and antibacterial activity of non-conventional curcuma species. *JBPNP* 9 (6), 457–464.
- Jain, A., Parihar, D.K., 2018. Antibacterial, biofilm dispersal and antibiofilm potential of alkaloids and flavonoids of Curcuma. *Biocatal Agric Biotechnol* 16, 677–682.
- Jain, A., Parihar, D.K., 2019. Molecular marker based genetic diversity study of wild, cultivated and endangered species of Curcuma from Chhattisgarh region for in situ conservation. *Biocatal Agric Biotechnol* 18, 101033.
- Jain, P., Pandey, R., Shukla, S.S., 2015. Acute and subacute toxicity studies of polyherbal formulation talisada churna in experimental animal model. *MJPMS* 1 (1), 7–10.
- Jain, P., Pandey, R., Shukla, S.S., 2016. Reproductive and developmental toxicity study of talisada churna: an ancient polyherbal formulation. *IAJPR* 6 (5), 5641–5653.
- Jain, P., Rao, S.P., Singh, V., Pandey, R., Shukla, S.S., 2014. Acute and sub-acute toxicity studies of an ancient ayurvedic formulation: agnimukha churna. *Columbia Journal of Pharmaceutical Sciences* (1), 18–22.
- Jain, P., Satapathy, T., Pandey, R.K., 2020 e e. A mini review of methods to control ticks population infesting cattle in Chhattisgarh with special emphasis on herbal acaricides. *IJNPR* 11 (12), 217–223.
- Jain, P., Satapathy, T., Pandey, R.K., 2020 c c. Efficacy of arecoline hydrobromide against cattle tick *Rhipicephalus (Boophilus) microplus*. *Int. J. Acarol* 46 (4), 268–275.
- Jain, P., Satapathy, T., Pandey, R.K., 2020 d d. First report on ticks (Acar: ixodidae) controlling activity of cottonseed oil (*Gossypium Sp.*). *Int. J. Acarol* 46 (4), 263–267.
- Jain, P., Satapathy, T., Pandey, R.K., 2020 a a. *Rhipicephalus microplus* (acari: ixodidae): clinical safety and potential control by topical application of cottonseed oil (*Gossypium sp.*) on cattle. *Exp. Parasitol.* 108017.
- Jain, P., Satapathy, T., Pandey, R.K., 2020 b b. *Rhipicephalus microplus*: a parasite threatening cattle health and consequences of herbal acaricides for upliftment of livelihood of cattle rearing communities in Chhattisgarh. *Biocatal Agric Biotechnol* 101611.
- Jain, S., Bhatia, G., Barik, R., Kumar, P., Jain, A., Dixit, V., 2009. Antidiabetic activity of *Paspalum scrobiculatum* Linn. in alloxan induced diabetic rats. *J. Ethnopharmacol.* 127, 325–328.
- Jaiswal, Y.S., Tatke, P.A., Gabhe, S.Y., Vaidya, A.B., 2016. Antidiabetic activity of extracts of *Anacardium occidentale* Linn. leaves on n-streptozotocin diabetic rats. *J Tradit Complement Med* 7 (4), 421–427.
- Janež, A., Guja, C., Mitrakou, A., Lalic, N., Tankova, T., Czupryniak, L., et al., 2020. Insulin therapy in adults with type 1 diabetes mellitus: a narrative review. *Diabetes*

- Ther 11 (2), 387–409.
- Joseph, B., Jini, D., 2013. Antidiabetic effects of *Momordica charantia* (bitter melon) and its medicinal potency. *Asian Pacific J Trop Dis* 3 (2), 93–102.
- Joseph, J., Joseph, M.G., Latha, S., 2018. Effect of aloe vera juice on glucose level among diabetics in a selected old age home at mangalore. 8 (9), 135–141.
- Joshi, B.C., Mukhija, M., Kalia, A.N., 2019. Phytochemistry and pharmacologic properties of *Urtica*. *Int. J. Green Pharm.* 201–209.
- Kanetkar, P., Singhal, R., Kamat, M., 2007. *Gymnema sylvestre*: a memoir. *J. Clin. Biochem. Nutr.* 41 (2), 77–81.
- Kaufman, N., Khurana, I., 2016. Using digital health technology to prevent and treat diabetes. *Diabetes Technol. Therapeut.* 18, 56–68.
- Khan, F., Sarker, M.M.R., Ming, L.C., Mohamed, I.N., Zhao, C., Sheikh, B.Y., et al., 2019. Comprehensive review on phytochemicals, pharmacological and clinical potentials of *gymnema sylvestre*. *Front. Pharmacol.* 10, 1223.
- Kianbakht, S., Hajiaghaei, R., 2013. Anti-hyperglycemic effects of *vaccinium arctostaphylos* L. fruit and leaf extracts in alloxan-induced diabetic rats. *J Med Plants* 12, 43–50.
- Kiran, P., Denni, M., Daniel, M., Laboratories, D., Farm, O.H., 2014. In: *Antidiabetic Principles , Phospholipids and Fixed Oil of Kodo Millet (Paspalum Scrobiculatum Linn.)*, vols. 13–5.
- Klangjareonchai, T., Roongpisuthipong, C., 2012. The effect of *tinospora crispa* on serum glucose and insulin levels in patients with type 2 diabetes mellitus. In: Shibata, M.-A. (Ed.), *J. Biomed. Biotechnol.* 2012, 808762.
- Krishna, R.N., Anitha, R., Ezhilarasan, D., 2020. Aqueous extract of *Tamarindus indica* fruit pulp exhibits antihyperglycaemic activity. *Avicenna J phytomedicine* 10 (5), 440–447.
- Kuczmánnová, A., Balažová, A., Račanská, E., Kameníková, M., Fialová, S., Majerník, J., Nagy, M., Gál, P., Mučají, P., 2016. *Agrimonia eupatoria* L. and *cynara cardunculus* L. Water infusions: comparison of anti-diabetic activities. *Molecules* 21 (5), 564.
- Kumar, V., Jain, P., Rathore, K., Ahmed, Z., 2016. Biological evaluation of pupalia lappacea for antidiabetic, antiadipogenic, and hypolipidemic activity both in vitro and in vivo. *Sci. Tech. Rep.* 9.
- Kumar, V., Rathore, K., Jain, P., Ahmed, Z., 2017. Biological activity of *bauhinia racemose* against diabetes and interlinked disorders like obesity and hyperlipidemia. *Clin. Phytoscience* 3, 7.
- Kumar, V., Van Staden, J., 2016. A review of *swertia chirayita* (gentianaceae) as a traditional medicinal plant. *Front. Pharmacol.* 6, 308.
- Kurup, S.B., 2017. Protective potential of *Averrhoa bilimbi* fruits in ameliorating the hepatic key enzymes in streptozotocin-induced diabetic rats. *Biomed. Pharmacother.* 85, 725–732.
- Kusano, S., Abe, H., 2000. Antidiabetic activity of white skinned sweet potato (*Ipomoea batatas* L.) in obese Zucker fatty rats. *Biol. Pharm. Bull.* 23 (1), 23–26.
- Latha, M., Pari, L., Ramkumar, K.M., Rajaguru, P., Suresh, T., Dhanabal, T., et al., 2009. Antidiabetic effects of scoparin acid D isolated from *Scoparia dulcis* in rats with streptozotocin-induced diabetes. *Nat. Prod. Res.* 23 (16), 1528–1540.
- Lawal, O.A., Ogunwande, I.A., Ibiroga, A.E., Layode, O.M., Opoku, A.R., 2015. Chemical constituents of essential oils from *catharanthus roseus* (L.) G. Don grown in Nigeria. *J. Essent Oil Bear Plants* 18 (1), 57–63.
- Leem, K.-H., Kim, M.-G., Hahn, Y.-T., Kim, H.K., 2016. Hypoglycemic effect of *opuntia ficus-indica* var. *sabotii* is due to enhanced peripheral glucose uptake through activation of AMPK/p38 MAPK pathway. *Nutrients* 8 (12), 800.
- Li, B., Cheng, M., Gao, H., et al., 2008. Back-regulation of six oxidative stress proteins with grape seed proanthocyanidin extracts in rat diabetic nephropathy. *J. Cell. Biochem.* 104, 668–679.
- Li, X., Yang, P., 2018. Research progress of *Sonchus* species Research progress of *Sonchus* species. *Int. J. Food Prop.* 21 (1), 147–157.
- Lindstedt, I., 2018. Ginger and diabetes : a mini-review. *Arch Gen Intern Med* 2 (2), 29–33.
- lodha, S.R., Joshi, S.V., Vyas, B.A., Upadhye, M.C., Kirve, M.S., Salunke, S.S., et al., 2010. Assessment of the antidiabetic potential of *Cassia grandis* using an *in vivo* model. "J. Adv. Pharm. Technol. Research" (JAPTR) 1 (3), 330–333.
- Luo, C., Zhang, W., Sheng, C., Zheng, C., Yao, J., Miao, Z., 2010. Chemical composition and antidiabetic activity of *opuntia milpa alta* extracts. *Chem. Biodivers.* 7, 2869–2879.
- Macédo, N.S., Silveira, Z. de S., Bezerra, A.H., Costa, J.G.M.da, Coutinho, H.D.M., Romano, B., et al., 2020. *Caesalpinia ferrea* C. Mart. (Fabaceae) phytochemistry, ethnobotany, and bioactivities: a review. *Molecules* 25 (17).
- Mahmoud, M.F., El Ashry, F.E., El Maraghy, N.N., Fahmy, A., 2017. Studies on the antidiabetic activities of *Momordica charantia* fruit juice in streptozotocin-induced diabetic rats. *Pharm. Biol.* 55 (1), 758–765.
- Mahmud, I., Shahria, N., Yeasmin, S., Iqbal, A., Mukul, E.H., Gain, S., Shilpi, J.A., Islam, M.K., 2018. Ethnomedicinal, phytochemical and pharmacological profile of a mangrove plant *Ceriops Decandra Griff*Din Hou. *J. Compl. Integr. Med.* 16 (1).
- Malarvili, A., Selvaraja, P., Ndyebura, A.W., Akowuh, G., Okechukwu, P., 2011. Antidiabetic activity of crude stem extracts of *coscinium fenestratum* on streptozotocin-induced type-2 diabetic rats. *Asian J. Pharmaceut. Clin. Res.* 4, 47–51.
- Mao, X., Wu, L.-F., Guo, H.-L., Chen, W.-J., Cui, Y.-P., Qi, Q., et al., 2016. The genus *Phyllanthus*: an ethnopharmacological, phytochemical, and pharmacological review. 2016. *Evid Based Complement Alternat Med*, 7584952.
- Mathew, J.J., Sajeshkumar, N.K., Joy, N., 2017. Phytochemical Characterization of *Averrhoa Bilimbi* and *In Vitro* Analysis of Cholesterol Lowering Effect on Fatty Food Materials. GRIN, Munich.
- McGaw, L.J., Srivastava, A.K., Lin, C.H., Steenkamp, V., 2019. Book Review: Medicinal Plants for Holistic Healing 10 p. 1053 *Frontiers in Pharmacology*.
- Medagama, A.B., 2015. The glycaemic outcomes of Cinnamon, a review of the experimental evidence and clinical trials. *Nutr. J.* 14 (1), 108.
- Meenakshi, P., Bhuvaneshwari, R., Rathi, M.A., Thirumoothi, L., Guravaiah, D.C., Jiji, M. J., et al., 2010. Antidiabetic activity of ethanolic extract of *zaleya decandra* in alloxan-induced diabetic rats. *Appl. Biochem. Biotechnol.* 162 (4), 1153–1159.
- Meloche, R.M., 2007. Transplantation for the treatment of type 1 diabetes. *World J. Gastroenterol.* 13 (47), 6347–6355.
- Mininel, F.J., Leonardo Junior, C.S., Espanha, L.G., Resende, F.A., Varanda, E.A., Leite, C. Q.F., et al., 2014. Characterization and quantification of compounds in the hydroalcoholic extract of the leaves from *Terminalia catappa* linn. (Combretaceae) and their mutagenic activity. In: van Vuuren, S. (Ed.), *Evidence-Based Complement Altern Med*, vol. 2014. p. 676902.
- Moqbel, F.S., Naik, P.R., Najma, H.M., Selvaraj, S., 2011. Antidiabetic properties of *Hibiscus rosa sinensis* L. leaf extract fractions on nonobese diabetic (NOD) mouse. *Indian J. Exp. Biol.* 49 (1), 24–29.
- Muñiz-Ramirez, A., Perez, R.M., Garcia, E., Garcia, F.E., 2020. Antidiabetic activity of aloe vera leaves. In: Sherman, R. (Ed.), *Evidence-Based Complement Altern Med*, vol. 2020. p. 6371201.
- Mwiti, K.C., Jide, A.A., 2015. Herbal therapy: a review of emerging pharmacological tools in the management of diabetes mellitus in Africa. 11 (44), 258–274.
- Nabi, S.A., Kasetti, R.B., Sirasanagandla, S., Tilak, T.K., Kumar, M.V.J., Rao, C.A., 2013. Antidiabetic and antihyperlipidemic activity of *Piper longum* root aqueous extract in STZ induced diabetic rats. *BMC Compl. Alternative Med.* 13, 37.
- Nayak, A., Sajini, J., Padmavati, D., Karthik, R., 2012. Chemistry and medicinal properties of *Coscinium fenestratum* (Gaertn.) colebr (tree turmeric). *Res. J. Pharm. Technol.* 5, 198–202.
- Nazreen, S., Kaur, G., Alam, M.M., et al., 2011. Hypoglycemic activity of *Bambusa arundinacea* leaf ethanolic extract in streptozotocin induced diabetic rats. *Pharmacologyonline* 1, 964–972.
- Niamat, R., Khan, M.A., Khan kiran, Y., Ahmed, M., Mazari, P., Ali, B., et al., 2012. A review on *zizyphus* as antidiabetic. *J. Appl. Pharmaceut. Sci.* 2, 177–179.
- Obih, P.O., Obih, C.A., Brown, A., Day, X., 2017. In vitro investigations of the antidiabetic action of ginger (*Zingiber officinale*). *Faseb. J.* 31 (1) 347–347.
- Okokon, J., Umoh, E., Etim, E., Jackson, C., 2009. Antiplasmoidal and antidiabetic activities of ethanol leaf extract of *heinsia crinata*. *J. Med. Food* 12, 131–136.
- Olivas-Quintero, S., López-Angulo, G., Montes-Avila, J., Díaz-Camacho, S.P., Vega-Aviña, R., López-Valenzuela, J.A., et al., 2017. Chemical composition and biological activities of *Helicteres vegae* and *Heliopsis sinaloensis*. *Pharm. Biol.* 55 (1), 1473–1482.
- Oram, R.A., Sims, E.K., Evans-Molina, C., 2019. Beta cells in type 1 diabetes: mass and function; sleeping or dead?. *Diabetologia* 62 (4), 567–577.
- Pal, R., Bhadada, S.K., Misra, A., 2021. COVID-19 vaccination in patients with diabetes mellitus: current concepts, uncertainties and challenges. *Diabetes Metab. Syndr.* 15 (2), 505–508.
- Palhetka, I.C., Ferreira, L.R., 2018. Hypoglycemic potential of *Anacardium occidentale* L. *J. Anal Pharm Res* 7 (2), 152–153.
- Pandit, R., Phadke, A., Jagtap, A., 2010. Antidiabetic effect of *Ficus religiosa* extract in streptozotocin-induced diabetic rats. *J. Ethnopharmacol.* 128 (2), 462–466 Mar 24.
- Paramanick, D., Sharma, N., 2017. A review on herbs which are used in diabetes mellitus. *International Journal of Drug Development* 9 (3), 12–17.
- Parsaeyan, N., 2012. The effect of coriander seed powder consumption on atherosclerotic and cardioprotective indices of type 2 diabetic patients. *Iran J Diabetes Obes* 4, 86–90.
- Peixoto Araujo, N.M., Arruda, H.S., de Paulo Farias, D., Molina, G., Pereira, G.A., Pastore, G.M., 2021. Plants from the genus *Eugenia* as promising therapeutic agents for the management of diabetes mellitus: a review. *Food Res. Int.* 142, 110182.
- Peter, K., Gandhi, P., 2017. Rediscovering the therapeutic potential of *Amaranthus* species: a review. *Egypt J Basic Appl Sci* 4 (3), 196–205.
- Prada, A.L., Amado, J.R.R., Keita, H., Zapata, E.P., Carvalho, H., Lima, E.S., et al., 2018. *Cassia grandis* fruit extract reduces the blood glucose level in alloxan-induced diabetic rats. *Biomed. Pharmacother.* 103, 421–428.
- Prakasam, A., Sethupathy, S., Pugalendi, K.V., 2005. Influence of *Casearia esculenta* root extract on glycoprotein components in streptozotocin diabetic rats. *Pharmazie* 60 (3), 229–232.
- Rahmani, A.H., 2015. *Cassia fistula* Linn: potential candidate in the health management. *Pharmacogn. Res.* 7 (3), 217–224.
- Raina, R., Verma, P.K., Peshin, R., Kour, H., 2019. Potential of *Juniperus communis* L as a nutraceutical in human and veterinary medicine. *Heliyon* 5 (8), e02376.
- Rajendran, M.P., Pallaiyan, B.B., Selvaraj, N., 2014. Chemical composition, antibacterial and antioxidant profile of essential oil from *Murraya koenigii* (L.) leaves. *Avicenna J phytomedicine* 4 (3), 200–214.
- Rao, A.R., Veeresham, C., Asres, K., 2013. In vitro and *in vivo* inhibitory activities of four Indian medicinal plant extracts and their major components on rat aldose reductase and generation of advanced glycation endproducts. *Phytother. Res.* 27, 753–760.
- Rao, S.P., Amrit, I., Jain, P., Singh, V., 2014. Antilulcer activity of agnimukha churna. *Int. J. Ayur. Pharma Research.* 2 (2), 40–46.
- Rao, S.P., Amrit, I., Singh, V., Jain, P., 2015. Antilulcer activity of natural compounds: a review. *Res. Rev. J. Pharmacogn. Phytochem.* 7 (2), 124–130.
- Rao, S.P., Jain, P., Rathore, P., Singh, V.K., 2016. Larvicidal and knockdown activity of *Citrus limetta* Risso oil against dengue virus vector. *IJNPR* 7 (3), 256–260.
- Rasineni, K., Bellamkonda, R., Singareddy, S.R., Desireddy, S., 2010. Antihyperglycemic activity of *Carthamus roseus* leaf powder in streptozotocin-induced diabetic rats. *Pharmacogn. Res.* 2 (3), 195–201.
- Rathore, K., Singh, V., Jain, P., Rao, S.P., Ahmed, Z., Thakur, V.S., 2014. In- vitro and *In vivo* antidiadipogenic, antidiabetic and hypolipidemic activity of *Diospyros melanoxylon* (Roxb.). *J. Ethnopharmacol.* 155, 1171–1176.
- Rathore, P., Rao, S.P., Roy, A., Satapathy, T., Singh, V., Jain, P., 2014. Hepatoprotective activity of isolated herbal compounds. *Res. J. Pharm. Technol.* 7 (2).

- Rivera-Mondragón, A., Ortíz, O.O., Bijttebier, S., Vlietinck, A., Apers, S., Pieters, L., et al., 2017. Selection of chemical markers for the quality control of medicinal plants of the genus *Cecropia*. *Pharm. Biol.* 55 (1), 1500–1512.
- Saeedi, P., Petersohn, I., Salpea, P., Malanda, B., Karuranga, S., Unwin, N., et al., 2019. Global and regional diabetes prevalence estimates for 2019 and projections for 2030 and 2045: results from the international diabetes federation diabetes atlas. ninth ed. 157. *Diabetes Res Clin Pract*.
- Salehi, B., Ata, A., Anil Kumar N, V., Sharopov, F., Ramírez-Alarcón, K., Ruiz-Ortega, A., et al., 2019. Antidiabetic potential of medicinal plants and their active components. *Biomolecules* 9 (10).
- Satyanarayana, K., Sravanthi, K., Shaker, I.A., Ponnulakshmi, R., 2015. Molecular approach to identify antidiabetic potential of *Azadirachta indica*. *J. Ayurveda Integr. Med.* 6 (3), 165–174.
- Sebei, K., Sakouhi, F., Herchi, W., Khouja, M.L., Boukhchina, S., 2015. Chemical composition and antibacterial activities of seven *Eucalyptus* species essential oils leaves. *Biol. Res.* 48 (1), 7.
- Senthilkumar, G., Arulselvan, P., Kumar, D., Subramanian, S., 2006. Anti-diabetic activity of fruits of *Terminalia chebula* on streptozotocin induced diabetic rats. *J. Health Sci.* 52, 283–291.
- Sepici-Dincel, A., Acikgoz, S., Cevik, C., et al., 2007. Effects of *in vivo* antioxidant enzyme activities of myrtle oil in normoglycaemic and alloxan diabetic rabbits. *J. Ethnopharmacol.* 110, 498–503.
- Shabani, E., Sayemiri, K., Mohammadpour, M., 2019. The effect of garlic on lipid profile and glucose parameters in diabetic patients: a systematic review and meta-analysis. *Prim Care Diabetes* 13 (1), 28–42.
- Sharwan, G., Jain, P., Pandey, R., Shukla, S.S., 2016. Toxicity and safety profiles of methanolic extract of *pistacia integerrima* J. L. Stewart ex brandis (PI) for wistar rats. *J. Pharmacopuncture* 19 (3), 253–258.
- Sharwan, G., Jain, P., Pandey, R., Shukla, S.S., 2015. Toxicity profile of traditional herbal medicine. *J. Ayu Herb Med* 1 (3), 81–90.
- Shi, C., Karim, S., Wang, C., Zhao, M., Murata, G., 2014. A review on antidiabetic activity of *Citrullus colocynthis* Schrad. *Acta Pol. Pharm.* 71 (3), 363–367.
- Shirwaikar, A., Rajendran, K., Dinesh Kumar, C., Bodla, R., 2004. Antidiabetic activity of aqueous leaf extract of *Annona squamosa* in streptozotocin-nicotinamide type 2 diabetic rats. *J. Ethnopharmacol.* 91 (1), 171–175.
- Shukla, S., Gunjegaonkar, S.M., 2018. Pharmacognostical and pharmacological profiling of *Barleria*. *J. Biol. Sci. Med.* 4, 41–50.
- Sibiya, A.M., Ramya, A.K., Vaseeharan, B., 2020. Bioactive molecules from *Phyllanthus niruri* and investigating their effects against diabetes. *Recent Trends in Biochemistry* 3, 1–7.
- Singh, A., 2018. Navneet. Critical review on various ethnomedicinal and pharmacological aspects of *piper longum* linn. (long pepper or pippali). *LJIPSR* 6 (1), 48–60.
- Singh, J., Cumming, E., Manoharan, G., Kalasz, H., Adeghate, E., 2011. Medicinal chemistry of the anti-diabetic effects of *Momordica charantia*: active constituents and modes of actions. *Open Med Chem* 5 (2), 70–77.
- Singh, P., Jain, P., Pandey, P., Shukla, S.S., 2016. Phytotherapeutic review on diabetes. *Spatula* 66 (2).
- Sinha, D., 2019. Ethnobotanical and pharmacological importance of western himalayan fir *abies pindrow* (Royle ex D. Don) Royle : A Review 31 (6), 1–14.
- Sk, A., Chakrapani, C., Alasyam, N., 2017. Evaluation of antidiabetic effect of *Murraya koenigii* leaves chloroform extract (MKLCE) in alloxan induced diabetic albino rats. *Pharma Innov.* J 6, 474–477.
- Skyler, J.S., Bakris, G.L., Bonifacio, E., Darsow, T., Eckel, R.H., Groop, L., et al., 2017. Differentiation of diabetes by pathophysiology, natural history, and prognosis. *Diabetes* 66 (2), 241 LP – 255.
- Su, J., Jiang, J., Zhang, F., Liu, Y., Ding, L., Chen, S., et al., 2019. Current achievements and future prospects in the genetic breeding of chrysanthemum: a review. *Hortic. Res.* 6 (1), 109.
- Tanaka, M., Misawa, E., Ito, Y., Habara, N., Nomaguchi, K., Yamada, M., et al., 2006. Identification of five phytosterols from aloe vera gel as antidiabetic compounds. *Biol. Pharm. Bull.* 29, 1418–1422.
- Teimouri, M., 2015. The chemical composition and antimicrobial activity of essential oils of *vaccinium arctostaphylos* L. *Int. J. Adv. Biol. Biom. Res* 3 (3), 270–274.
- Teugwa, C.M., Mejia, P.C., Zofou, D., Tchinda, B.T., Boyom, F.F., 2013. Antioxidant and antidiabetic profiles of two African medicinal plants: *picalima nitida* (Apocynaceae) and *Sonchus oleraceus* (Asteraceae). *BMC Compl. Alternative Med.* 13, 175.
- Tiwari, P., Mishra, B.N., Sangwan, N.S., 2014. Phytochemical and pharmacological properties of *gymnema sylvestre*: an important medicinal plant. In: Vincent, J.B. (Ed.), *BioMed Res. Int.* 2014, 830285.
- Verma, D.S., Vashishth, E., Singh, R., Kumari, A., Meena, A., Pant, P., et al., 2013. A review on parts of *Albizia lebbeck* (L.) Benth. used as Ayurvedic drugs. *Res. J. Pharm. Technol.* 6, 1307–1313.
- Verma, S., Gupta, M., Popli, H., Aggarwal, G., 2018. Diabetes Mellitus Treatment Using Herbal Drugs 10 (1), 1–10.
- Vilar, D. de A., Vilar, M.S. de A., de Lima e Moura, T.F.A., Raffin, F.N., de Oliveira, M.R., Franco, C.F. de O., et al., 2014. Traditional uses, chemical constituents, and biological activities of *Bixa orellana* L.: a review. *ScientificWorldJournal* 2014, 857292.
- Wang, B., Chou, K.H., Queenan, B.N., Pennathur, S., Bazan, G.C., 2019. Molecular design of a new diboronic acid for the electrohydrodynamic monitoring of glucose. *Angew. Chem.* 58 (31), 10612–10615.
- Wang, C., Yang, J., Zhao, P., Zhou, Q., Mei, Z., Yang, G., et al., 2014. Chemical constituents from *Eucalyptus citriodora* Hook leaves and their glucose transporter 4 translocation activities. *Bioorg. Med. Chem. Lett* 24 (14), 3096–3099.
- Wani, S.A., Fenugreek, K.P., 2018. A review on its nutraceutical properties and utilization in various food products. *J. Saudi Soc Agric Sci* 17 (2), 97–106.
- WHO. World Health Organization. mHealth, 2011. New Horizons for Health through Mobile Technologies: Second Global Survey on eHealth.
- Xu, Y., Liang, J.Y., 2005. Chemical constituents of *Sonchus oleraceus* L. *J. China Pharm. Univ.* 36, 411–413.
- Yakoob, A.T., Tajuddin, N.B., Hussain, M.I.M., Mathew, S., Govindaraju, A., Qadri, I., 2016. Antioxidant and hypoglycemic activities of *clausena anisata* (wildd.) hook f. Ex benth. Root mediated synthesized silver nanoparticles. *Pharmacogn J*, 589–586.
- Yamamoto, J., Tadaishi, M., Yamane, T., Oishi, Y., Shimizu, M., Kobayashi-Hattori, K., 2015. Hot water extracts of edible *Chrysanthemum morifolium* Ramat. exert antidiabetic effects in obese diabetic KK-Ay mice. *Biosci. Biotechnol. Biochem.* 79 (7), 1147–1154.
- Yan, Y.M., Fang, P., Yang, M.T., Li, N., Lu, Q., Cheng, Y.X., 2015. Anti-diabetic nephropathy compounds from *Cinnamomum cassia*. *J. Ethnopharmacol.* 165, 141–147.
- Yang, Y.C., Mong, M.C., Wu, W.T., Wang, Z.H., Yin, M.C., 2020. Phytochemical profiles and anti-diabetic benefits of two edible *Amaranthus* species. *CyTA - J. Food* 18 (1), 94–101.
- Yasir, M., Das, S., Kharya, M.D., 2010. The phytochemical and pharmacological profile of *Persea americana* Mill. *Pharmacogn Rev* 4 (7), 77–84.
- Yobe, M., Togenu, F.B., Joyce, O., Augustine, U., 2017. Research article gas chromatographic characterization of aqueous saponin. Extract of *Heinsia crinata* Leaves 9 (8), 201–203.
- Yu, H., Zhang, P., Liu, H., Sun, X., Liang, J., Sun, L., et al., 2021. Hypoglycemic activity of *Origanum vulgare* L. and its main chemical constituents identified with HPLC-ESI-QTOF-MS. *Food Funct.* 12, 2580–2590.
- Zahid, M., Mujahid, M., Singh, P.K., Farooqui, S., Singh, K., Parveen, S., Arif, M., 2018. *Annona squamosa* Linn. (custard apple): an aromatic medicinal plant fruit with immense nutraceutical and therapeutic potentials. *Int. J. Pharma Sci. Res.* 9 (5), 1745–1759.
- Zaid, H., Mahdi, A.A., Tamrakar, A.K., Saad, B., Razzaque, M.S., Dasgupta, A., 2016. Natural active ingredients for diabetes and metabolism disorders treatment. *Evidence-Based Complement Altern Med.* 2016, 2965214.
- Zengin, G., Locatelli, M., Stefanucci, A., Macedonio, G., Novellino, E., Mirzaie, S., et al., 2017. Chemical characterization, antioxidant properties, anti-inflammatory activity, and enzyme inhibition of *Ipomoea batatas* L. leaf extracts. *Int. J. Food Prop.* 20 (2), 1907–1919.
- Zheng, M.S., Liu, Y.S., Yuan, T., Liu, L.Y., Li, Z.Y., Huang, X.L., 2020. [Research progress on chemical constituents of *Citrullus colocynthis* and their pharmacological effects]. *Zhongguo Zhongyao Zazhi* 45 (4), 816–824.